arXiv:1111.4662v3 [math.PR] 6 Jun 2012

The distributions of traffics and their free product: an asymptotic freeness theorem for random matrices and a central limit theorem

Camille Male*

ABSTRACT:

The distributions of traffics are defined and are applied for families of larges random matrices, random groups and infinite random rooted graphs with uniformly bounded degree. There are constructed by adding axioms in Voiculescu's definition of *-distribution of non commutative random variables. The convergence in distribution of traffics generalizes Benjamini, Schramm, Aldous, Lyons' weak local convergence of random graphs. We introduce a notion of freeness of traffics, which contains both the classical notion of independence and Voiculescu's notion of freeness. We prove an asymptotic freeness theorem for families of matrices invariant by permutation, which enlarges the class of large random matrices for which we can predict the empirical eigenvalues distribution. We prove a central limit theorem for the sum of free traffics, and interpret the limit as the (traffic)-convolution of a gaussian commutative random variable and a semicircular non commutative random variable. We make a connection between the freeness of traffics and the natural free product of random graphs, combination of the statistical independence and of the geometric free product.

Overview of the article

0.1 Motivation from random matrix theory

In all this article, when we consider an N by N matrix H_N, we implicitly mean that we consider a sequence of matrices $(H_N)_{N\geqslant 1}$, such that H_N is N by N. Following the random matrix theory terminology, the empirical eigenvalues distribution of an N by N matrix H_N is the probability measure

$$\mathcal{L}_{H_N} = \frac{1}{N}\sum_{i=1}^{N} \delta_{\lambda_i}, \tag{0.1}$$

where $\lambda_1, \dots, \lambda_N$ are the eigenvalues of H_N and δ_λ is the Dirac mass in λ. The convergence of empirical eigenvalues distributions considered in this article is the convergence in moments, i.e. the convergence of

$$\mathcal{L}_{H_N}(P) = \frac{1}{N}\sum_{i=1}^{N} P(\lambda_i, \bar{\lambda}_i)$$

for any polynomial P. When H_N is random we consider the convergence in expectation: we say that in expectation H_N has a limiting eigenvalues distribution whenever $\mathbb{E}\big[\mathcal{L}_{H_N}(P)\big]$ converges for any polynomial P.

A simple question is the following.

---

*UMPA, ENS de Lyon, 46 allée d'Italie, 69007 Lyon, France. camille.male@ens-lyon.fr

Question 1 (The spectrum of the sum of two matrices).
Let A_N, B_N be two independent random N by N Hermitian matrices whose entries have all their moments. Let U_N be an N by N random unitary matrix, independent of (A_N, B_N) and distributed according to the uniform distribution on the set of permutation matrices. How can we predict the possible limit of the empirical eigenvalues distribution of the matrix

$$H_N = A_N + U_N B_N U_N^* \text{ ?} \tag{0.2}$$

Voiculescu [26] introduced free probability theory, which gives the concepts and the tools to analyze limiting eigenvalues distributions of random matrices. From its pioneering connection between large random matrices and freeness [26], Question 1 has been solved in many cases. To quote a few examples, the answer is known when

- B_N is invariant in law under conjugacy by a unitary matrix, A_N and B_N are uniformly bounded in operator norm [2],
- B_N is a Wigner matrix and A_N is uniformly bounded in operator norm [14] or is a structured matrix [4],
- B_N is a random covariant matrix and A_N is uniformly bounded in operator norm [10],
- A_N and B_N are distributed according to the Haar measure on the unitary group [2] or the permutation group [22, 21],
- A_N and B_N are band matrices [25], etc.

Consider such a random matrix

$$H_N = A_N + U_N B_N U_N^*$$

and assume that B_N is invariant under conjugacy by unitary matrices. Regarding Question 1, this is equivalent to assume that U_N is actually distributed according to the Haar measure on the unitary group. Voiculescu [27] introduced the notion of *-probability spaces, with an notion of convergence in *-distribution, and the notion of *-freeness, a non commutative analogue of the notion of statistical independence (see Sections 0.4 and 0.5 below). The algebra of random matrices with all their moments is a *-probability space and the description of the limiting eigenvalues distribution of H_N is given by Voiculescu's asymptotic freeness theorem.

This theorem implies the following result. Assume that

1. The empirical eigenvalues distribution of A_N (respectively B_N) converges to a probability measure $\mathcal{L}_a$ (respectively $\mathcal{L}_b$),
2. A_N and B_N are uniformly bounded in operator norm.

Then, H_N has limiting eigenvalues distribution which is the so-called free convolution of the limiting eigenvalues distributions of A_N and B_N:

$$\mathcal{L}_h = \mathcal{L}_a \boxplus \mathcal{L}_b.$$

It is clear that this formula is not valid for A_N and B_N arbitrary when U_N is a uniform permutation matrix instead of a Haar matrix. Indeed, assume that A_N and B_N are diagonal matrices with limiting eigenvalues distribution $\mathcal{L}_a$ and $\mathcal{L}_b$ respectively. Then, the empirical eigenvalues distribution of H_N converges to the classical convolution of $\mathcal{L}_a$ and $\mathcal{L}_b$:

$$\mathcal{L}_h = \mathcal{L}_a * \mathcal{L}_b.$$

Classical and free convolutions are known to be different operations in most of the cases. In a nutshell, we need more than the limiting spectra of A_N and B_N in order to predict the limiting spectrum of H_N (the *-distribution of a Hermitian matrix is its empirical eigenvalues distribution). Formalizing the information needed on A_N and B_N to infer the spectrum of H_N is the task of the notion of distribution of traffics.

0.2 Convergence in distribution of traffics for a single matrix

Let A_N be an N by N random complex matrix. We give a construction of the distribution of traffics of A_N starting by the observation: if we want to answer Question 1 in a general context, the notion of distribution we introduce must be invariant by conjugacy of A_N by permutation matrices. By a formal analysis of joint moments of random matrices invariant by permutation, we see that it will be convenient to introduce a certain set of graphs as "test functions".

Let U_N be a random permutation matrix, uniformly distributed and independent of A_N. Let

$$X_N = U_N A_N U_N^*$$

which is a random matrix invariant in law by permutation. The distribution of A_N must be the distribution of X_N.

Now, since we only consider convergence in expectation, the mean distribution of a random matrix must depend only on the joint law of its entries, say of its mixed moments. So, we first write the joint moments of the entries of X_N conditionally on A_N.

Let $K = (k_{i,j})_{i,j\geqslant 1}$ and $L = (\ell_{i,j})_{i,j\geqslant 1}$ be families of non negative integers, with finite support in $\{1,\dots,N\}^2$ (K and L are fixed whereas N will go to infinity). Denote the associated joint moments of the entries of X_N

$$\delta^0_{K,L}(A_N) = \mathbb{E}\Big[\prod_{i,j=1}^N X_N(i,j)^{k_{i,j}} \overline{X_N(i,j)}^{\ell_{i,j}} \Big| A_N \Big]. \tag{0.3}$$

By invariance by permutation of X_N, for any permutation σ of $\{1,\dots,N\}$,

$$\begin{aligned} \delta^0_{K,L}(A_N) &= \mathbb{E}\Big[\prod_{i,j=1}^N X_N\big(\sigma(i),\sigma(j)\big)^{k_{i,j}} \overline{X_N(\sigma(i),\sigma(j))}^{\ell_{i,j}} \Big| A_N \Big] & (0.4)\\ &= \mathbb{E}\Big[\prod_{i,j=1}^N X_N(i,j)^{k_{\sigma(i),\sigma(j)}} \overline{X_N(i,j)}^{\ell_{\sigma(i),\sigma(j)}} \Big| A_N \Big]. & (0.5) \end{aligned}$$

It turns out that the dependence on K,L in $\delta^0_{K,L}(A_N)$ can be expressed in terms of a finite graph $T_{K,L}$, with possibly loops and multiple edges, and whose edges are labelled by symbols a and a^* (see Figure 1). Consider first the graph on the vertices $1,\dots,N$, with $k_{i,j}$ directed edges labelled a and $\ell_{j,i}$ directed edges labelled a^* for any pair of vertices (i,j). Remove the vertices that are not attached to any edge, and call the resulting graph $T_{K,L}$. Hence, $\delta^0_{A_N}[T] = \delta^0_{K,L}(A_N)$ depends only of the isomorphism class T of the labelled graph $T_{K,L}$.

A finite graph with possibly loops and multiple edges, whose edges are labelled by symbols a and a^* is called a *-graph in one variable in this article. The set of finite connected *-graphs in one variable is denoted by $\mathcal{G}\langle x,x^*\rangle$. A element of this set is called a *-test graph. A *-test graph is said to be cyclic whenever there exists a paths visiting each edge once in the sense of its orientation. The set of cyclic *-test graphs is denoted by $\mathcal{G}_{cyc}\langle x,x^*\rangle$.



Figure 1: The *-graph which is associated to $\mathbb{E}\big[\overline{X_N(i_2,i_1)}X_N(i_2,i_2)^2\overline{X_N(i_3,i_2)}X_N(i_3,i_1)\big|A_N\big]$, with i_1,i_2,i_3 distinct. For a better readability, edges labelled a are represented by a simple arrow whereas edges labelled a^* are represented by an arrow with a cross.

Definition 0.1 (Injective trace and convergence in distribution of traffics)**.**
For any **-test graph* T *in* $\mathcal{G}\langle x, x^*\rangle$*, we set*

$$\tau^0_{A_N}[T] = \frac{(N-1)!}{(N-|V|)!}\delta^0_{A_N}[T], \tag{0.6}$$

where V *is the set of vertices of* T *and* $|V|$ *is its cardinal. The number* $\tau^0_{A_N}[T]$*, also denoted* $\tau^0_N\big[T(A_N)\big]$*, is called the injective trace of* T *taken in* A_N*. The convergence in distribution of traffics of a matrix* A_N *is the pointwise convergence of* $\tau^0_{A_N} : \mathcal{D} \to \mathbb{C}$*, where* $\mathcal{D}$ *is a subset of* **-test graphs, called the domain of convergence of* τ_{A_N}*,*

$$\mathcal{G}_{cyc}\langle x, x^*\rangle \subset \mathcal{D} \subset \mathcal{G}\langle x, x^*\rangle,$$

stable by identification of vertices (see Definition 3.4). If A_N *is random, we consider the convergence in expectation, and say that in expectation* A_N *has a limiting distribution of traffics whenever* $\mathbb{E}\big[\tau^0_{A_N}[T]\big]$ *converges for any* T *in* $\mathcal{D}$*.*

We usually denote by τ^0_a the limiting distribution of $\tau^0_{A_n}$, τ^0_b for $\tau^0_{B_N}$, etc. This notation is justified since we construct a model for a, called a traffic. The map τ^0_a can be viewed as a Fourier transform of a map $\tau_a : \mathcal{D} \to \mathbb{C}$ called the distribution of traffics of a (see Section II). This transform is invertible.

0.3 Convergence in distribution of traffics as a convergence of networks

In this article, we call network a directed graph, with no multiple edges nor loops, whose edges are labelled by random variables. A finite network can be encoded by a matrix when we chose an enumeration of its vertices. Such a matrix is called an adjacency matrix of the network. Since the distribution of traffics of a matrix is invariant under conjugacy by permutation matrices, the distribution of traffics of a finite network is well defined as the distribution of traffics of any of its adjacency matrices. Hence, Definition 0.1 gives a mode of convergence for large finite networks.

There exists many ways to define a continuous limit of a large graph or a large network, as the notions of graphons [17, 18] and graphings [6, 1, 3, 15] (see below). To compare these definitions, we interpret geometrically the injective trace for adjacency matrices of graphs. Let A_N be an adjacency matrix of a finite directed graph G_N with no multiple edges nor loops: for any $m, n = 1, \dots, N$,

$$A_N(m,n) = \left\{ \begin{array}{ll} 1 & \text{if } (n,m) \text{ is a vertex of } G_N \\ 0 & \qquad\text{otherwise.} \end{array} \right.$$

Let T be a test graph in one variable (a *-test graph with no edge labelled a^*). Denote its set of vertices by V. Then,

1. $\delta^0_{A_N}[T]$ is the probability that a random injective map $\Phi_N : V \to \{1, \dots, N\}$ is a graph morphism $T \to G_N$ (that is a map preserving adjacency of vertices and orientation).
2. $\tau^0_{A_N}[T]$ is the expectation of the number of root preserving injective morphisms $\Phi_N : T \to G_N$, where a root of T is chosen arbitrarily, and a root of G_N is chosen uniformly at random.

Convergence from the right and graphons: Following graph theory terminology [17], we call injective density profile of G the map $\delta^0_{G_N} : T \mapsto \delta^0_{A_N}[T]$, when T varies among the set of all test graphs. The pointwise convergence of the injective density profile of a graph is called the convergence form the right. A graph whose injective density profile converges has a continuous limiting model, called a graphons. This mode of convergence is applyed for dense graphs, i.e. graphs with an number of edges of order N^2, as for the complete graph. The injective density profile of a finite network is the map $T \mapsto \delta^0_{A_N}[T]$ defined with the same formula [8], the convergence from the right is the point-wise convergence of this map.

Weak local convergence and graphings: The convergence in distribution of traffics of graphs is, in a non straightforward way, the weak local convergence (see Part V). The later has been introduced by Benjamini and Schramm [6] in a specific case and extended by Aldous and Lyons [1]. The notion of graphing is a continuous model for weak local limits of graphs. The weak local convergence of a random network is usually defined as the weak local convergence of its graphs, together with the convergence in law of its random labels (conditionally on its graph). This differs from the distribution of traffics.

Very recently, the notion of graphlets [11] has been introduced as a continuous model of networks limit.

0.4 Distribution of traffics as a generalized *-distribution

We go back to the definition of distributions of traffics of random matrices. We introduce them in comparison with the *-distributions of free probability. Let $\mathbf{A}_N = (A_1, \dots, A_p)$ be a family of N by N matrices. The *-distribution of $\mathbf{A}_N$ is the map

$$\begin{array}{rccc} \Phi_{\mathbf{A}_N}: & \mathbb{C}\langle \mathbf{x}, \mathbf{x}^* \rangle & \to & \mathbb{C} \\ & P & \mapsto & \frac{1}{N}\mathrm{Tr}\big[P(\mathbf{A}_N, \mathbf{A}_N^*)\big]. \end{array} \tag{0.7}$$

The symbol $\mathbb{C}\langle \mathbf{x}, \mathbf{x}^* \rangle$ denotes the set of non commutative polynomials in indeterminates $\mathbf{x} = x_1, \dots, x_p$ and $\mathbf{x}^* = x_1^*, \dots, x_p^*$. The symbol Tr denotes the usual trace of N by N matrices, sum of diagonal elements. We use $P(\mathbf{A}_N, \mathbf{A}_N^*)$ as a shortcut for $P(A_1, \dots, A_p, A_1^*, \dots, A_p^*)$.

The convergence in *-distribution of $\mathbf{A}_N$ is the point-wise convergence of $\Phi_{\mathbf{A}_N}$. When $\mathbf{A}_N$ is a family of random matrices, we consider the convergence in expectation, that is the convergence of $\mathbb{E}\big[\Phi_{\mathbf{A}_N}(P)\big]$ for any polynomial P. If A_N is a normal matrix, denote by $\lambda_1, \dots, \lambda_N$ its eigenvalues. Then, for any polynomial P in two (commutative) indeterminates

$$\frac{1}{N}\mathrm{Tr}\big[P(A_N, A_N^*)\big] = \frac{1}{N}\sum_{i=1}^N P(\lambda_i, \bar\lambda_i) = \mathcal{L}_{A_N}(P),$$

where $\mathcal{L}_{A_N}$ denotes the empirical eigenvalues distribution of A_N. Hence, *-distributions of random matrices are a generalization of empirical eigenvalues distributions. For a normal matrix the knowledge of its *-distribution is equivalent to the knowledge of its empirical eigenvalues distribution.

We now write $\Phi_{\mathbf{A}_N}$ in terms of the entries of the matrices. We will see how defining the distributions of traffics of $\mathbf{A}_N$ by capturing more information on the matrices than for the *-distribution. The *-distribution of $\mathbf{A}_N$ is equivalent to the collection of all numbers $\Phi_{\mathbf{A}_N}(P)$, for any monic monomial P, that is all complex numbers

$$\frac{1}{N}\sum_{i_1, \dots, i_K = 1}^N A_{\gamma(1)}^{\varepsilon(1)}(i_1, i_2) A_{\gamma(2)}^{\varepsilon(2)}(i_2, i_3) \dots A_{\gamma(K-1)}^{\varepsilon(K-1)}(i_{K-1}, i_K) A_{\gamma(K)}^{\varepsilon(K)}(i_K, i_1), \tag{0.8}$$

for any integer $K \geqslant 1$, any map $\varepsilon : \{1, \dots, K\} \to \{1, *\}$ and any map $\gamma : \{1, \dots, K\} \to \{1, \dots, p\}$. Restricted on cyclic *-test graphs, the distribution of traffics of $\mathbf{A}_N$ is actually the collection of the numbers as in (0.8) where some of the indices are constraint to be equal. More generally, we set

$$m_{\pi, \varepsilon, \gamma}(\mathbf{A}_N) = \frac{1}{N}\sum_{\ker \mathbf{i} \geqslant \pi} A_{\gamma(1)}^{\varepsilon(1)}(i_1, i_2) A_{\gamma(2)}^{\varepsilon(2)}(i_3, i_4) \dots A_{\gamma(K)}^{\varepsilon(K)}(i_{2K-1}, i_{2K}), \tag{0.9}$$

where

- ε varies among the set of maps $\{1, \dots, K\} \to \{1, *\}$,
- γ varies among the set of maps $\{1, \dots, K\} \to \{1, \dots, p\}$,
- π varies among the set of partitions of $1, \dots, 2K$,

- and the sum is over all families of integers $\mathbf{i} = (i_1, \dots, i_{2K})$ in $\{1, \dots, N\}$ such that integers whose indices are in a same block of π are constraint to be equal: $n \sim_\pi m \Rightarrow i_n = i_m$, for all $n, m = 1, \dots, 2K$.

It turns out that the number $m_{\pi,\varepsilon,\gamma}(\mathbf{A}_N)$ depends only on $\mathbf{A}_N$ and on a *-graph in indeterminates $\mathbf{x}$: that is a finite graph with possibly loops and multiple edges, whose edges are labelled by symbols $x_1, \dots, x_p, x_1^*, \dots, x_p^*$, see Figure 2. Let $T_{\pi,\varepsilon,\gamma}$ be the graph whose vertices are the blocks of the partition π. For any $n, m = 1, \dots, 2K$, there is on directed edge in $T_{\pi,\varepsilon}$ from the block of n and the block of m for each occurrence of a term $A_\gamma^\varepsilon(i_n, j_m)$ in the product. Such an edge is labelled x_γ^ε.



Figure 2: The *-graph which is associated to $\frac{1}{N}\sum_{i_1,i_2,i_2} A_N^*(i_1,i_2)A_N(i_2,i_2)^2A_N^*(i_2,i_3)A_N(i_3,i_1)$. The graph is the same as in Figure 1, but the associated numbers are different, since in the sum i_1, i_2, i_3 may be equal.

The number $m_{\pi,\varepsilon,\gamma}(\mathbf{A}_N)$ depends only on the isomorphism class T of $T_{\pi,\varepsilon,\gamma}$. It is denoted by $\tau_{\mathbf{A}_N}[T]$ or $\tau_N\big[T(\mathbf{A}_N)\big]$, and is called the trace of T in $\mathbf{A}_N$. The distribution of traffics of $\mathbf{A}_N$ is the map

$$\begin{array}{cccc} \tau_{A_N}: & \mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle & \to & \mathbb{C} \\ & T & \mapsto & \tau_N\big[T(A_N)\big], \end{array} \tag{0.10}$$

where $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ denotes the set of *-test graphs in indeterminates $\mathbf{x}$, that is the set of finite, connected, finite graphs whose edges are labelled by indeterminates $x_1, \dots, x_p, x_1^*, \dots, x_p^*$. Let $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$ the the subset of $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ whose graphs possesses a cycle visiting each edge once in the sense of their orientation. The convergence in distribution of traffics of $\mathbf{A}_N$ is the pointwise convergence of $\tau_{\mathbf{A}_N}$ on a domain

$$\mathcal{G}_{cyc}\langle x, x^* \rangle \subset \mathcal{D} \subset \mathcal{G}\langle x, x^* \rangle,$$

stable by identification of vertices (see Definition 3.4).

The quantities (0.9) have been introduced by Mingo and Speicher in [20] (see Section 10). They construct operators associated to graphs whose edges are labelled by matrices (called representations of quivers). From their construction, we can interpret the numbers (0.9) as normalized traces of product of operators between tensor product of $\mathbb{C}^N$ (see Part III). We also interpret their main theorem in [20] as a result of sequential compactness in distribution of traffics.

A limiting distribution of traffics $\tau : \mathcal{D} \to \mathbb{C}$ induces a *-distribution $\Phi^{(\tau)}$ since $\mathcal{D}$ is assumed to contains cycles, that correspond to the numbers in (0.8). Even for a single normal matrix A_N, the distribution τ_{A_N} contains more than its empirical eigenvalues distribution $\mathcal{L}_{A_N}$.

We give a solution of Question 1, about characterizing the possible limiting spectrum of

$$H_N = A_N + U_N B_N U_N^*$$

in the context of distribution of traffics: we describe the mean limiting distribution of traffics of H_N in terms of the limiting distribution of traffics of A_N and B_N, their existence assumed. By the above, we then get the convergence in moments of the empirical eigenvalues distribution of H_N.

Given a domain $\mathcal{G}_{cyc}\langle x, x^* \rangle \subset \mathcal{D} \subset \mathcal{G}\langle x, x^* \rangle$, stable by identification of vertices, and a map

$\tau : \mathcal{D} \to \mathbb{C}$ that satisfies a property of positivity, we construct a *-probability space: that is a *-algebra endowed with a positive linear form, still denoted $\Phi^{(\tau)}$, such that $\Phi^{(\tau)}(\mathbf{1}) = 1$. Elements of such a space are called traffics. Question 1 about the spectrum of H_N is reformulated in terms of the spectrum of the traffic

$$h = a + b$$

where a and b are traffics constructed as in the next Section.

0.5 The freeness of traffics, asymptotic freeness

The freeness of distribution of traffics is defined as the rule that emerges naturally from the following theorem.

Theorem 0.2 (The asymptotic freeness of $(A_N, U_N B_N U_N^*)$).
Let A_N and B_N be two deterministic matrices and let U_N be a uniform permutation matrix. Assume that A_N (respectively B_N) has a limiting distribution of traffics τ_a (respectively τ_b) on $\mathcal{G}\langle x, x^*\rangle$. Then, in expectation $(A_N, U_N B_N U_N^*)$ has a limiting distribution of traffics $\tau_{(a,b)}$ described as follow. Consider a *-test graph T whose edges are labelled by variables a, a^*, b and b^*, obtained as a "free product" of *-test graphs in a, a^* and in b, b^* only (see Figure 3 and Section 11). Then $\tau^0_{(a,b)}[T]$ is the product of the injective traces τ^0_a or τ^0_b of the connected components of T in a same (and corresponding) color. If T is not such a free product, then $\tau^0_{(a,b)}[T] = 0$.



Figure 3: A free product of *-graphs in indeterminates a and a^* (red) and b and b^* (black). Subgraphs attached to vertices of a given colored component are distinct.

Two traffics a and b whose joint distribution is as in Theorem 0.2 are said to be free (see Part IV). The freeness for distribution of traffics does not always implies the *-freeness in the sense of Voiculescu of the induced *-distribution $\Phi^{(\tau)}$.

Recall the definition of *-freeness (see [2]). The sub-algebras $\mathcal{A}_1, \dots, \mathcal{A}_p$ of a *-probability space $(\mathcal{A}, .^*, \Phi)$ are free if and only if $\Phi(a_j) = 0, a_j \in \mathcal{A}_{i_j}$ and $i_j \neq i_{j+1}$ for all $j \geqslant 1$ implies $\Phi(a_1 \dots a_n) = 0$ for all $n \geqslant 1$. Non commutative random variables $a_1, \dots, a_p$ are said to be free whenever the algebras they span are *-free.

The distribution $\Phi^{(\tau)}$ induced by a distribution of traffics τ depends only on the knowledge of τ in a small domain (on *-test graphs that are colored cycles). How to determine when the freeness of traffics induces the *-freeness of the non commutative random variables is not a problem tackled in this article.

0.6 Between *-freeness and independence

Considered again the two situations of the first Section:

- A_N^s and B_N^s are unitarily invariant Hermitian matrices,
- A_N^d and B_N^d are diagonal real matrices.

Let U_N be a uniform permutation matrix, independent of $(A_N^s, B_N^s, A_N^d, B_N^d)$. Then, under technical assumptions, Theorem 12.1 gives the convergence of the Hermitian matrices

$$H_N^x = A_N^x + U_N B_N^x U_N^*, \quad x \in \{s, d\},$$

and its limit it characterized by the freeness of the limiting distribution of traffics of A_N^x and B_N^x. By Voiculescu's asymptotic freeness theorem [2], the limiting eigenvalues distribution of H_N^s is given by the free convolution of the limiting eigenvalues distributions of A_N^s and B_N^s. By a straightforward computation, the limiting eigenvalues distribution of H_N^d is given by the classical convolution of the limiting eigenvalues distributions of A_N^d and B_N^d. Hence, the freeness of traffics encodes simultaneously the classical notion of statistical independence of random variables and Voiculescu's *-freeness (see Proposition 11.2 and Corollary 12.7) of non commutative random variables. Seeing the freeness of traffics as a mixture of statistical independence and Voiculescu's freeness is also suggested by the two following results.

We prove a central limit theorem for the sum of free traffics which "interpolates" Laplace and Voiculescu's central limit theorems.

Theorem 0.3 (A central limit theorem for the sum of traffics)**.**
Let $t_1, \dots, t_n$ be identically distributed, centered, self adjoint, free traffics. Then, the traffic

$$m_n = \frac{t_1 + \dots + t_n}{\sqrt{n}}$$

converges in distribution of traffics on the domain of cyclic *-test graphs to the sum $d + s$ of free traffics, where d is the "traffic-incarnation" of a Gaussian commutative random variable and s the "traffic-incarnation" of a semicircular non commutative random variables.

Furthermore, following Benjamini, Schramm, Aldous, Lyons approach [6, 1], one can define the distribution of traffics of rooted random graphs with uniformly bounded degree. For such random graphs, we can define a natural notion of free products, where each copies of components of the free product are sampled independently each other (see Figure 4 and Section 16.1).

Theorem 0.4 (The free products of random graphs)**.**
Let G_1 and G_2 be free independent, traffic-stationary, integrable, rooted random graphs with uniformly bounded degree. Then the distribution of the sum $H = G_1 + G_2$ where G_1 and G_2 are realized as traffics corresponds to the random free product of G_1 and G_2.

0.7 Applications

Theorem 12.1 implies the convergence in moments of the mean empirical eigenvalues distribution for any normal random matrix H_N which can be written as

$$H_N = P(\mathbf{X}_N, \mathbf{U}_N, \mathbf{U}_N^*, \mathbf{G}_N, \mathbf{Y}_N, \mathbf{Y}_N^*), \tag{0.11}$$

where P is a fixed non commutative polynomial (which does not depend on N) and

1. $\mathbf{X}_N = (X_1^{(N)}, \dots, X_p^{(N)})$ is a family of independent random N by N real or complex Wigner matrices (Corollary 12.7).
2. $\mathbf{U}_N = (U_1^{(N)}, \dots, U_q^{(N)})$ is a family of independent real or complex uniform permutation matrices (Corollary 12.9).
3. $\mathbf{G}_N = (G_1^{(N)}, \dots, G_r^{(N)})$ is a family of independent adjacency matrices of random graphs with N vertices, permutation invariant, having a so-called weak local limit (see Part V).



Figure 4: A realization of the free product with itself of the random graph which is $\mathbb{Z}$ with probability 0.8, and the graph with one vertex and no edge with probability 0.2 (truncated at distance 10 of the root).

4. $\mathbf{Y}_N = (Y_1^{(N)}, \dots, Y_s^{(N)})$ is a family of arbitrary N by N deterministic matrices, having a limiting distribution of traffics (Corollary 12.5).
5. $\mathbf{X}_N, \mathbf{U}_N, \mathbf{G}_N$ being independent.

It also implies the convergence of certain observables of eigenvectors of H_N when H_N is Hermitian (see Propositions 2.2 and 12.6). Moreover, these convergences hold actually for a larger class of matrices than polynomials in the matrices above-named (see Sections 5 and 8). The explicit computation of such limiting eigenvalues distributions remains a hard task and is not tackled in this article.

An other application of the asymptotic freeness theorem of this article is the following (see Section 12.2.1). For any unitary matrix U_N having a limit u in distribution of traffics, one can naturally associate an operation between probability measures with compact support. This operation could be called u-convolution. Extremal cases are classical and free convolutions. The family of such operations is much bigger than the interpolation in [5].

Organization of the paper:
In Part I, we introduce the notion of distribution of traffics and their convergence for a family of matrices. We show how we can extract information about the macroscopic behavior of random matrices from their distribution of traffics. In part II, we introduce the injective trace and gives examples of limiting distribution of traffics. In part III, we define general distribution of traffics

and space of traffics. We also translate a result of Mingo and Speicher [20] in terms of distributions of traffics. In part IV, we introduce the notion of freeness, state a result of asymptotic freeness for random matrices and a central limit theorem for the sum of free traffics. In part V, we prove the connection between convergence in the sense of traffics and weak local convergence of graphs, and free products of traffics and free products of random graphs.

Acknowledgment:
The author would like to gratefully thank Mikael de la Salle , Alice Guionnet, Florent Benaych-Georges and Charles Bordenave for useful discussions. He also acknowledges the financial support of the ANR GranMa.

This paper has known significant progress during some travels in 2011: a visit of Ashkan Nikeghbali and Kim Dang at the Institut für Mathematik of Zurich, the school ”Vicious walkers and random matrices“ in les Houches organized by the CNRS, and the summer school ”Random matrix theory and its applications to high-dimensional statistics“ in Changchun, founded jointly by the CNRS of France and the NSF of China. The author gratefully acknowledges the organizers of these events for providing an inspiring working environment.

Part I

The distribution of traffics of a family of matrices

Notation for graphs

A (directed) graph (with possible multiple edges) G is a couple (V, E), where V is a set, referred as the set of vertices of G, and E is a multi-set, referred as the set of edges of G. Recall that a multi-set is a generalization of the notion of set in which members are allowed to appear more than once. When E is finite, it is denoted

$$E = \{\!\!\{ \underbrace{e_1, \dots, e_1}_{k_1}, \dots, \underbrace{e_p, \dots, e_p}_{k_p} \}\!\!\}, \tag{0.12}$$

where for any $j = 1, \dots, p$, e_j is an element of V^2. Formula (0.12) means that for any $j = 1, \dots, p$, the element e_j appears exactly k_j times in E. This number is called the multiplicity of e_j. The graph G is said to be finite when both V and E are finite.

Graphs are always considered up to isomorphisms: two graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are isomorphic whenever there exists a bijection $\phi : V_1 \to V_2$ preserving the adjacency of vertices, their orientation and their multiplicity.

1 Definition

1.1 Graphs in several variables

Definition 1.1 (Graphs in several variables).

1. *A graph in p variables is a graph whose edges are labelled by free variables $x_1, \dots, x_n$. Formally, a graph in p variables consists in a triplet $T = (V, E, \gamma)$ where (V, E) is a graph and γ is a map $E \to \{1, \dots, p\}$ which indicates the indeterminate corresponding to each edge.*
2. *A *-graph in p variables is a graph in $2p$ free variables $x_1, \dots, x_p$, $x_1^*, \dots, x_p^*$. Formally, it consists in a quadruple $T = (V, E, \gamma, \varepsilon)$, where (V, E, γ) is a test graph in p variables and ε is a map $E \to \{1, *\}$, which indicates which indeterminates are transposed.*

The vertices and the edges of $T = (V, E, \gamma)$ refer to the vertices and the edges of the graph (V, E). For a graph $T = (V, E, \gamma)$ in one variable, the map γ is trivial and then T is identified with (V, E). A *-graph in one variable is simply denoted $T = (V, E, \varepsilon)$. Some examples of *-graphs are drawn in Figure 5.

Remark about the Figures:
For conciseness, in all the figures of the article, we use colors instead of labels when illustrating *-graphs in several variables. Moreover, when the edges labelled x_i are plotted in a certain color, the edges labelled x_i^* are plotted with the same color with a cross.

1.2 A generalization of the trace

Now we define the observables we extract from a family of matrices.

Definition 1.2 (The trace of a *-graph in a family of matrices).
For any family $\mathbf{A}_N = (A_1, \dots, A_p)$ of $N \times N$ complex matrices and any finite *-graph $T = (V, E, \gamma, \varepsilon)$ in p variables we set

$$\tau_N\big[T(\mathbf{A}_N)\big] := \frac{1}{N} \sum_{\phi : V \to \{1, \dots, N\}} \prod_{e \in E} A_{\gamma(e)}^{\varepsilon(e)}\big(\phi(e)\big), \tag{1.1}$$



Figure 5: Six *-test graphs in three variables, a (red), b (black) and c (blue).

where

- *for any directed edge* $e = (v_1, v_2)$, *we have set* $\phi(e) = (\phi(v_1), \phi(v_2))$,
- *and for any* N *by* N *matrix* M *and any integers* n, m *in* $\{1, \dots, N\}$, *the complex number* $M(n, m)$ *is the entry* (n, m) *of* M.
- M^* *is the conjugate transpose of the matrix* M

We say that $\tau_N\big[T(\mathbf{A}_N)\big]$ *is the trace of the* **-graph* T *in the matrices* $\mathbf{A}_N$.

Examples: We explicit the quantities that correspond to the first four *-graphs drawn Figure 5, from left to right. Let $\mathbf{A}_N = (A_N, B_N, C_N)$ be a family of N by N matrices. We use the convention that in any *-graph T of Figure 5, the red edges stand for A_N, the black ones stand for B_N and the blue ones stand for C_N.

a) $\tau_N\big[T(\mathbf{A}_N)\big] = \frac{1}{N}\sum_{i=1}^N A_N(i,i) = \frac{1}{N}\mathrm{Tr}[A_N]$, where Tr stands for the trace of matrices, i.e. the sum of diagonal elements.

b) $\tau_N\big[T(\mathbf{A}_N)\big] = \frac{1}{N}\sum_{i,j=1}^N A_N(i,j)B_N^*(j,i) = \frac{1}{N}\mathrm{Tr}[A_N B_N^*]$.

c) $\tau_N\big[T(\mathbf{A}_N)\big] = \frac{1}{N}\sum_{i,j,k=1}^N B_N(i,j)B_N(j,k)A_N(k,i) = \frac{1}{N}\mathrm{Tr}[B_N B_N^t A_N]$, where $\cdot^t$ stands for the transpose.

d) $\tau_N\big[T(\mathbf{A}_N)\big] = \frac{1}{N}\sum_{i=1}^N A_N(i,i)B_N(i,i) = \frac{1}{N}\mathrm{Tr}[A_N \circ B_N]$, where $\circ$ stands for the Hadamard product of matrices, i.e. the entry-wise product.

Remark that if T is a finite *-graph with connected components $T_1, \dots, T_K$, one has

$$\tau_N\big[T(\mathbf{A}_N)\big] = N^{K-1}\tau_N\big[T_1(\mathbf{A}_N)\big]\dots\tau_N\big[T_K(\mathbf{A}_N)\big].$$

1.3 The distribution of traffics of a family of matrices

Definition 1.3 (*-test graphs)**.**

1. *A test graph is a finite, connected graph in several variables.*
2. *A* **-test graph is a finite, connected* **-graph in several variables.*
3. *A test graph or a* **-test graph is said to be cyclic whenever there exists a cycle on its graph visiting each edge exactly once, in the sense of their orientation.*

The set of all test graphs in p variables is denoted by $\mathcal{G}\langle x_1, \dots, x_p\rangle$, where the symbols $x_1, \dots, x_p$ refer to the indeterminates. We use the shortcut $\mathcal{G}\langle \mathbf{x}\rangle = \mathcal{G}\langle x_1, \dots, x_p\rangle$ in the sequel, the number of variables being implicit. The set of all *-test graphs in p variables is denoted in short by $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle$. The set of all cyclic test graphs in p variables is denoted by $\mathcal{G}_{cyc}\langle \mathbf{x}\rangle$, the set of all cyclic *-test graphs is denoted by $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^*\rangle$. In Figure 5, all but the second and the rightmost *-test graphs are cyclic.

Definition 1.4 (Formal distribution of traffics)**.** *A formal distribution of traffics is a map $\tau : \mathcal{D} \to \mathbb{C}$, where*

$$\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle \subset \mathcal{D} \subset \mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle.$$

The set $\mathcal{D}$ is called the domain of τ.

Definition 1.5 (Distribution of traffics of a family of matrices)**.** *The distribution of traffics of a family $\mathbf{A}_N = (A_1^{(N)}, \dots, A_p^{(N)})$ of $N \times N$ complex matrices is the map*

$$\begin{array}{cccc} \tau_{\mathbf{A}_N} : & \mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle & \to & \mathbb{C} \\ & T & \mapsto & \tau_{\mathbf{A}_N}[T] := \tau_N\big[T(\mathbf{A}_N)\big]. \end{array}$$

Remarks about symmetries:

- The distribution of traffics of a family $\mathbf{A}_N = (A_1, \dots, A_p)$ of N by N complex matrices is invariant under joint conjugacy by any permutation matrix: let U_N be a permutation matrix associated to a permutation σ of $\{1, \dots, N\}$, that is, for any $i, j = 1, \dots, N$,

  $$U_N(i,j) = \left\{ \begin{array}{ll} 1 & \text{if } \sigma(i) = j \\ 0 & \text{otherwise.} \end{array} \right. \tag{1.2}$$

  Then the distribution of traffics of $\mathbf{A}_N$ is the same as the distribution of the jointly conjugated family

  $$U_N \mathbf{A}_N U_N^* = (U_N A_1 U_N^*, \dots, U_N A_p U_N^*). \tag{1.3}$$

- If the matrices of the family $\mathbf{A}_N$ are symmetric, then for any *-test graph T the number $\tau_N\big[T(\mathbf{A}_N)\big]$ does not depend on the direction of the edges of T.

- If the matrices are Hermitian, the number $\tau_N\big[T(\mathbf{A}_N)\big]$ is equal to $\tau_N\big[\tilde{T}(\mathbf{A}_N)\big]$, where $\tilde{T}$ is the test graph obtained by replacing labels x_k^* by x_k for any k

- If the matrices of the family $\mathbf{A}_N$ are real, then $\tau_N\big[T(\mathbf{A}_N)\big]$ is equal to $\tau_N\big[\tilde{T}(\mathbf{A}_N)\big]$, where $\tilde{T}$ is the test graph obtained by reversing edges labelled x_k^* and replacing this label by x_k for any k.

- If the matrices of the family $\mathbf{A}_N$ are $0-1$ matrices, i.e. their entries are either 0 either 1, then $\tau_N\big[T(\mathbf{A}_N)\big]$ is equal to $\tau_N\big[\tilde{T}(\mathbf{A}_N)\big]$, where $\tilde{T}$ is the *-test graph obtained by forgetting the multiplicity of edges with the same label.

1.4 Convergence in distribution of traffics

From now, when we consider a family $\mathbf{A}_N$ of complex matrices we implicitly mean that we have considered a sequence $(\mathbf{A}_N)_{N \geqslant 1}$ of p-tuples for a certain p, where for any $N \geqslant 1$ the matrices in $\mathbf{A}_N$ are N by N.

Definition 1.6 (Convergence in distribution of traffics)**.**
Let $\mathbf{A}_N$ be a family of complex matrices and $\mathcal{D}$ be a domain

$$\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle \subset \mathcal{D} \subset \mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle.$$

We say that $\mathbf{A}_N$ has a limiting distribution of traffics τ on $\mathcal{D}$ whenever $\tau_{\mathbf{A}_N}$ converges pointwise on $\mathcal{D}$ to τ, i.e.

$$\forall T \in \mathcal{D}, \quad \tau[T] = \lim_{N \to \infty} \tau\big[T(\mathbf{A}_N)\big] \quad \textit{exists.} \tag{1.4}$$

If the matrices of $\mathbf{A}_N$ are random, we say that in expectation $\mathbf{A}_N$ has a limiting distribution of traffics τ on $\mathcal{D}$ whenever

$$\forall T \in \mathcal{D}, \quad \tau[T] = \lim_{N \to \infty} \mathbb{E}\Big[\tau\big[T(\mathbf{A}_N)\big]\Big] \quad \textit{exists.} \tag{1.5}$$

Many usual matrix models has a limiting distribution of traffics with domain the full set of *-test graphs $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ (see Section 4). The set $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$ is the minimal subset of $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ for which we can use the method of injective trace to recover the limiting spectrum of large matrices (see Section 3).

An other domain is introduced in Section 10. The set of proper *-test graphs $\mathcal{G}_{pr}\langle \mathbf{x}, \mathbf{x}^* \rangle$ which is a maximal domain of sequential compactness with respect to convergence in distribution of traffics.

In the following, the domain of convergence will always be $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ or $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$. Even when we have the convergence on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ for a given matrix model, we sometimes focus on the limiting distribution restricted on $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$ since it happens than this notion is more universal than the later. For instance, real and complex Wigner matrices has the same limiting distribution of traffics on $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$, be not in $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ (see Proposition 4.2).

2 Eigen-structure of large random matrices in their limiting distribution of traffics

Given a normal matrix having a limiting distribution of traffics τ, we can compute from τ many features about its limiting eigen-structure. The convergence of the empirical eigenvalues distribution is automatic (see Proposition 2.1). We also get limiting observables of eigenvectors of a matrix from its distribution of traffics τ, as it is illustrated in Section 2.2.

2.1 Limiting empirical eigenvalues distribution

Let H_N be an N by N normal matrix. The empirical eigenvalues distribution of H_N is probability measure on $\mathbb{C}$

$$\mathcal{L}_{H_N} = \frac{1}{N} \sum_{i=1}^{N} \delta_{\lambda_i}, \tag{2.1}$$

where $\lambda_1, \dots, \lambda_N$ are the eigenvalues of H_N and δ_λ is the Dirac mass in λ.

Proposition 2.1 (Convergence of the empirical eigenvalues distribution)**.** *If H_N has a limiting distribution of traffics, then its empirical eigenvalues distribution converges in moments.*

Proof. The convergence in moments of $\mathcal{L}_{H_N}$ is the convergence for any monic monomial P in two commutative indeterminates of

$$\mathcal{L}_{H_N}(P) := \frac{1}{N} \sum_{i=1}^{N} P(\lambda_i, \bar{\lambda}_i) = \frac{1}{N} \mathrm{Tr}\big[P(H_N, H_N^*) \big]. \tag{2.2}$$

If $P(z) = z^n \bar{z}^m$, then the later quantity is equal to $\tau_N\big[T(H_N, H_N^*)\big]$, where T is the cyclic *-test graph in one variable with vertices $1, \dots, n+m$ and edges $(1,2), (2,3), \dots, (n+m-1, n+m), (n+m, 1)$, n edges being labelled with the first variable and the other ones by the second (see Figure 6, picture on the left). Hence, $\mathcal{L}_{H_N}(P)$ converges as N goes to infinity. □



Figure 6: For the *-test graph T_1 on the left, $\tau_N\big[T_1(H_N)\big] = \tau_N[H_N^2 H_N^{*2}]$. For the one T_2 on the right, $\tau_N\big[T_2(H_N)\big] = \tau_N[H_N^3 \circ H_N^2]$.

2.2 A limiting observable of eigenvectors

Let H_N be an N by N Hermitian matrix. Since H_N is diagonalizable in a unitary basis, we can write

$$H_N = \sum_{k=1}^N \lambda_k u_k u_k^*, \tag{2.3}$$

where $\lambda_1 \leqslant \ldots \leqslant \lambda_N$ are the eigenvalues of H_N and $u_1, \ldots, u_N$ in $\mathbb{C}^N$ are associated eigenvectors.

Proposition 2.2 (Eigenvectors and distribution of traffics).
If H_N has a limiting distribution of traffics on $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$, then for any polynomials P and Q

$$\Delta(H_N) = \frac{1}{N} \sum_{m,n=1}^N P(\lambda_m) Q(\lambda_n) \sum_{i=1}^N \big(|u_n(i)|^2 \ |u_m(i)|^2 - \frac{1}{N^2} \big) \tag{2.4}$$

converges as N goes to infinity.

Proof. For any real functions f and g,

$$\begin{aligned} \frac{1}{N}\mathrm{Tr}\big[f(H_N) \circ g(H_N) \big] &= \frac{1}{N} \sum_{i=1}^N \big(f(H_N) \big)(i,i) \big(g(H_N) \big)(i,i) \\ &= \frac{1}{N} \sum_{n,m=1}^N f(\lambda_n) g(\lambda_m) \sum_{i=1}^N |u_n(i)|^2 |u_m(i)|^2. \end{aligned}$$

Recall that $\circ$ stands for the Hadamard product of matrices, and that by functional calculus,

$$f(H_N) := \sum_{k=1}^N f(\lambda_k) u_k u_k^*.$$

Hence,

$$\begin{aligned} &\frac{1}{N}\mathrm{Tr}\big[f(H_N) \circ g(H_N) \big] - \frac{1}{N}\mathrm{Tr}\big[f(H_N) \big] \frac{1}{N}\mathrm{Tr}\big[g(H_N) \big] \\ &= \frac{1}{N} \sum_{n,m=1}^N f(\lambda_n) g(\lambda_m) \sum_{i=1}^N \big(|u_n(i)|^2 |u_m(i)|^2 - \frac{1}{N^2} \big) \end{aligned}$$

If f and g are polynomials, then the quantities $\frac{1}{N}\mathrm{Tr}\big[f(H_N) \circ g(H_N) \big]$, $\frac{1}{N}\mathrm{Tr}\big[f(H_N) \big]$ and $\frac{1}{N}\mathrm{Tr}\big[g(H_N) \big]$ converges as N goes to infinity. This is known for the normalized trace by the previous section. By multi-linearity its is enough to prove that $\frac{1}{N}\mathrm{Tr}\big[f(H_N) \circ g(H_N) \big]$ converges when f and g are monic monomials. If $f(x) = x^n$ and $g(x) = x^m$, then this quantity is equal to $\tau_N\big[T(H_N) \big]$ where T is the cyclic test graph in one variable with vertices $1, \ldots, n, 1', \ldots, m'$ and edges $(1,2), (2,3), \ldots, (n-1, n), (n,1)$ and $(1', 2'), (2', 3'), \ldots, (m'-1, m'), (m', 1)$ (see Figure 6, picture on the right). □

Part II

The injective trace, examples of limiting distributions of traffics

The injective version of a formal distribution of traffics can heuristically be seen as its cumulant version, or as its Fourier transform. It is a tool for the computation of limiting distribution of traffics of random matrices invariant under permutation and for Hadamard product of matrices (see Wigner matrices and complex permutation matrices described below). It is also involved in the definition of freeness for traffics (see Part IV).

3 The injective trace

3.1 Definition

Definition 3.1 (The injective trace for families of matrices).
Let $T = (V, E, \gamma, \varepsilon)$ be a *-graph in p variables and $\mathbf{A}_N = (A_1, \dots, A_p)$ be a family of N by N matrices. The injective trace of T in $\mathbf{A}_N$ is

$$\tau_N^0\big[T(\mathbf{A}_N)\big] = \frac{1}{N} \sum_{\substack{\phi:V\to\{1,\dots,N\}\\ injective}} \prod_{e\in E} A_{\gamma(e)}^{\varepsilon(e)}\big(\phi(e)\big).$$

By the inclusion-exclusion principle, we can write the injective trace of a *-graph in a family of matrices as a weighted sum of the trace of *-graphs in the matrices, and reciprocally. To write the relation between τ_N and τ_N^0 properly, we need the following definition (see Figure 7).

Given a *-graph $T = (V, E, \gamma, \varepsilon)$ and a partition π of V, we define a new *-graph $\pi(T) = \big(\pi(V), \pi(E), \pi(\gamma), \pi(\varepsilon)\big)$, where we have identified the vertices that belong to a same block. The set of vertices $\pi(V)$ are the blocks of π. If V is the multiset $\big\{\!\!\big\{(v_1, v_2), \dots, (v_{2K-1}, v_{2K})\big\}\!\!\big\}$, then $\pi(V)$ is the multiset $\big\{\!\!\big\{(\pi(v_1), \pi(v_2)), \dots, (\pi(v_{2K-1}), \pi(v_{2K}))\big\}\!\!\big\}$, where for any v in V, $\pi(v)$ denotes the block of π containing v. For any $e = (\pi(v_{2k-1}), \pi(v_{2k}))$ in $\pi(V)$, we set $\pi(\gamma)(e) = \gamma(v_{2k-1}, v_{2k})$ and $\pi(\varepsilon)(e) = \varepsilon(v_{2k-1}, v_{2k})$.

Lemma 3.2 (Injective trace vs. non-injective trace).
Let $T = (V, E, \gamma)$ be a *-graph in p variables. Then, for any p-tuple $\mathbf{A}_N$ of $N \times N$ matrices, one has

$$\tau_N\big[T(\mathbf{A}_N)\big] = \sum_{\pi\in\mathcal{P}(V)} \tau_N^0\big[\pi(T)(\mathbf{A}_N)\big], \tag{3.1}$$

where $\mathcal{P}(V)$ is the set of partitions of V. Hence, one has

$$\tau_N^0\big[T(\mathbf{A}_N)\big] = \sum_{\pi\in\mathcal{P}(V)} \tau_N\big[\pi(T)(\mathbf{A}_N)\big] \times \mu_V(\pi), \tag{3.2}$$

where μ_V is the Möbius function of the finite poset $\mathcal{P}(V)$ (see [23]*).*

3.2 Convergence in distribution of traffics

Let $\mathcal{D}$ be a domain

$$\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^*\rangle \subset \mathcal{D} \subset \mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle.$$

Assume that $\mathcal{D}$ is stable by "identification of vertices", i.e. for any T in $\mathcal{D}$ and π partition of the vertices of T, one has $\pi(T) \in \mathcal{D}$. This property holds for the set $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^*\rangle$ of cyclic *-test graphs ($\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^*\rangle$ is actually the smallest class of *-test graphs stable by identification of vertices and possessing cycles as the leftmost example of Figure 6).

Then, the following proposition is an immediate consequence of Lemma 3.2.



Figure 7: Starting with the leftmost test graph T, this Figure represents all the test graphs $\pi(T)$ obtained for any partition π of the vertices of T (which includes T itself). Hence, the trace of T is the sum of the injective trace of all the test graphs in the Figure.

Proposition 3.3 (Convergence in injective trace)**.**
A family $\mathbf{A}_N$ *of matrices has a limiting distribution of traffics* τ *with domain* $\mathcal{D}$ *if and only if* $\tau^0_N\big[T(\mathbf{A}_N)\big]$ *converges for any* T *in* $\mathcal{D}$*.*

This motivates the following definition.

Definition 3.4 (Injective version of a formal distribution of traffics)**.**
Let $\tau : \mathcal{D} \to \mathbb{C}$ *be a formal distribution of traffics. Assume that its domain* $\mathcal{D}$ *is stable by identification of vertices. The injective version of* τ *is the map* $\tau^0 : \mathcal{D} \to \mathbb{C}$ *defined by: for any* **-test graph* $T \in \mathcal{D}$*,*

$$\tau^0\big[T\big] \quad = \quad \sum_{\pi \in \mathcal{P}(V)} \tau\big[\pi(T)\big] \times \mu_V(\pi), \tag{3.3}$$

where μ_V *is as in Lemma 3.2. Hence, we have*

$$\tau\big[T\big] \quad = \quad \sum_{\pi \in \mathcal{P}(V)} \tau^0\big[\pi(T)\big]. \tag{3.4}$$

4 Examples of limiting distributions of traffics

4.1 Wigner matrices

Definition 4.1 (Wigner matrices)**.**
A random matrix X_N *is an* N *by* N *real or complex Wigner matrix whenever:*

1. *Almost surely,* X_N *is Hermitian, i.e.* $X_N = X_N^*$*,*
2. *The sub-diagonal entries of* X_N *are independent, identically distributed,*
3. *The diagonal entries of* $M_N = \sqrt{N} X_N$ *are distributed according to a measure* ν *on* $\mathbb{R}$*,*
4. *Real case: the strictly sub-diagonal entries of* M_N *are distributed according to a measure* μ *on* $\mathbb{R}$*,*

4'. *Complex case: for any* $k < l$, $M_N(k,l) = \frac{x_{k,l} + i y_{k,l}}{\sqrt{2}}$*, where* $x_{k,l}$ *and* $y_{k,l}$ *are independent, identically distributed according to a symmetric probability measure* μ *on* $\mathbb{R}$*,*

5. μ *and* ν *do not depend of* N *and possess all their moments.*

A real or complex Wigner matrix is almost surely Hermitian, so its distribution of traffics is completely characterized in $\mathcal{G}\langle x \rangle$.

Proposition 4.2 (The limiting distribution of a Wigner matrix)**.**
Let X_N *be an* N *by* N *real or complex Wigner matrix. Assume that* μ *and* ν *are centered and* μ *is of variance one, that is*

$$\int x d\mu(x) = \int x d\nu(x) = 0, \int x^2 d\mu(x) = 1.$$

Then, in expectation X_N *has a limiting distribution of traffics* τ *on* $\mathcal{G}\langle x \rangle$ *which does not depend on* μ *and* ν*. For any test graph* T *in one variables:*

- *Real case:* $\tau^0[T] = 1$ *if* T *is a double tree.*
- *Complex case:* $\tau^0[T] = 1$ *if* T *is a double tree whose twin edges have different directions.*
- $\tau^0[T] = 0$ *otherwise.*

Being a double tree for T *means that it is a tree when we forget the multiplicity and the orientation of its edges, and all (undirected) edges are of multiplicity two. Twin edges of a double tree is a pair of edges sharing the same vertices (see Figure 8).*

It should be noticed that a cyclic double tree has necessarily twin edges of different directions. Hence, real and complex Wigner matrices of Propositions 4.2 have the same limiting distribution of traffics on $\mathcal{G}_{cyc}\langle x\rangle$, but not in $\mathcal{G}\langle x\rangle$.



Figure 8: Two double trees. The leftmost does not contribute for complex Wigner matrices, since certain twin edges have the same orientation. It is non cyclic. The rightmost does and is cyclic.

Proof. Generalities about random matrices invariant by permutation:
Let M_N be an N by N random matrix with complex entries. We say that M_N is invariant by permutation if and only if for any permutation matrix U_N, one has

$$M_N \overset{\mathcal{L}}{=} U_N M_N U_N^*, \tag{4.1}$$

where the equality is in law of random matrices. If U_N is the matrix associated to the permutation σ, then the entry (i,j) of $U_N M_N U_N^*$ is $M_N\big(\sigma(i),\sigma(j)\big)$. For such matrices, the use of the injective trace makes the computation of the mean distribution of traffics straightforward. Indeed, for any *-test graph $T=(V,E,\varepsilon)$ in one variable,

$$\begin{aligned}\mathbb{E}\Big[\tau_N^0\big[T(M_N)\big]\Big] &= \mathbb{E}\Bigg[\frac{1}{N}\sum_{\substack{\phi:V\to\{1,\dots,N\}\\ \text{injective}}}\prod_{e\in E} M_N^{\varepsilon(e)}\big(\phi(e)\big)\Bigg]\\ &= \frac{1}{N}\sum_{\substack{\phi:V\to\{1,\dots,N\}\\ \text{injective}}}\mathbb{E}\Bigg[\prod_{e\in E} M_N^{\varepsilon(e)}\big(\phi(e)\big)\Bigg].\end{aligned}$$

But the quantity $\delta_N^0\big[T(M_N)\big] = \mathbb{E}\Big[\prod_{e\in E} M_N^{\varepsilon(e)}\big(\phi(e)\big)\Big]$ does not depend on the injective map ϕ. Hence, we get

$$\mathbb{E}\Big[\tau_N^0\big[T(M_N)\big]\Big] = \frac{(N-1)!}{(N-|V|)!}\delta_N^0\big[T(M_N)\big], \tag{4.2}$$

Assume that $\delta_N^0\big[T(M_N)\big]$ converges to a complex number $\delta^0[T]$. For instance, when the entries of M_N are independent identically distributed according to a law that does not depend on N, then $\delta_N^0\big[T(M_N)\big]$ does not depend on N. If we rescale the matrix, say by a factor N^α for an $\alpha>0$, we get

$$\mathbb{E}\Bigg[\tau_N^0\Big[T\Big(\frac{1}{N^\alpha}M_N\Big)\Big]\Bigg] = \big(N^{|V|-(\alpha|E|+1)}+o(1)\big)\delta^0\big[T\big].$$

Hence, *-test graphs with a small number of edges relatively to the number of vertices are in better position to contribute. Recall the following classical result of graph theory (see [16, Lemma 1.1] for a proof).

Lemma 4.3 (Number of edges and vertices in a connected graph).
Let $G = (V, E)$ be a finite connected graph. Then, one has

$$|V| \leqslant |E| + 1, \tag{4.3}$$

with equality if and only if G is a tree.

Proof of Proposition 4.2: the real case.
Consider a test graph $T = (V, E)$ in one variable x. The matrix $M_N = \sqrt{N} X_N$ is invariant under permutation and has independent identically distributed entries, so

$$\mathbb{E}\Big[\tau_N^0\big[T(X_N)\big]\Big] \quad = \quad N^{|V| - \left(\frac{|E|}{2}+1\right)} \times \delta^0(T) + o(1), \tag{4.4}$$

where $\delta^0[T] = \mathbb{E}\Big[\prod_{e \in E} M_N\big(\phi(e)\big)\Big]$ for any injective map $\phi : V \to \{1, \dots, N\}$. This quantity is independent of N.

The sub-diagonal entries of M_N are independent and centered and M_N is Hermitian. So $\delta^0[T]$ vanishes as soon as one (undirected) edge of T is of multiplicity one. Hence, using Lemma 4.3, we get that $\mathbb{E}\big[\tau_N^0\big[T(X_N)\big]\big]$ converges to zero except possibly if each edge is of multiplicity strictly bigger than one and $|V| = \frac{|E|}{2} + 1$. By the second part of Lemma 4.3, T is necessarily a double tree.

Since $\mathbb{E}\big[M_N(k,l)^2\big] = 1$ for any $k \neq l$, we get that for any double tree T,

$$\mathbb{E}\big[\tau_N^0\big[T(X_N)\big]\big] \underset{N \to \infty}{\longrightarrow} 1.$$

Proof of Proposition 4.2: the complex case.
Without any modification of the proof of the real case, we get that $\tau_N^0\big[T(X_N)\big]$ tends to zero if T is not a double tree.

Since $\mathbb{E}\big[M_N(k,l)^2\big] = 0$ and $\mathbb{E}\big[|M_N(k,l)|^2\big] = 1$ for any $k \neq l$, we get for any double tree T, $\mathbb{E}\big[\tau_N^0\big[T(X_N)\big]\big]$ converges to one if the twin edges of T are is the opposite directions, and to zero otherwise.

□

4.2 Uniform permutation matrices

Definition 4.4 (Real and complex uniform permutation matrices).
A random matrix U_N is an N by N real or complex uniform permutation matrix whenever for any $m, n = 1, \dots, N$,

$$U_N(m,n) = \left\{ \begin{array}{cc} 1 & \textit{if } \sigma_N(m) = n \textit{ (real case)} \\ e^{i\theta_{m,n}} & \textit{if } \sigma_N(m) = n \textit{ (complex case)} \\ 0 & \textit{otherwise,} \end{array} \right. \tag{4.5}$$

where

1. *σ_N is a random permutation on $\{1, \dots, N\}$ uniformly distributed,*
2. *$(\theta_{m,n})_{m,n=1,\dots,N}$ is a family of independent random variables uniformly distributed on $[0, 2\pi]$, independent of σ_N.*

A real uniform permutation matrix is called simply a uniform permutation matrix.

Proposition 4.5 (The limiting distribution of a permutation matrix)**.**
Let U_N be a real or complex uniform permutation matrix. Then, in expectation U_N has a limiting distribution of traffic τ on $\mathcal{G}\langle x\rangle$ given by: for any *-test graph T in one variable,

- *Real case: $\tau^0[T] = 1$ if T is a directed line,*
- *Complex case: $\tau^0[T] = 1$ if T is a Hermitian directed line,*
- *$\tau^0[T] = 0$ otherwise.*

Being a directed line for T means that there exists an integer $K \geqslant 1$ such that the vertices of T are $1, \dots, K$ and its directed edges are $(1,2), \dots, (K-1,K)$ labelled x and $(2,1), \dots, (K,K-1)$ labelled x^*, with arbitrary multiplicity. Being a Hermitian directed line for T means that the multiplicity of edges $(n,n+1)$ equals the multiplicity of edges $(n+1,n)$ for any $n = 1, \dots, K-1$ (see Figure 9).



Figure 9: Two directed lines. The leftmost does not contribute for complex uniform permutation matrices, since it is not a Hermitian *-test graph. It is non cyclic. The rightmost does and is cyclic.

As for real and complex Wigner matrices, it should be noticed that real and complex uniform permutation matrices have in expectation the same limiting distribution of traffics on cyclic test graphs.

Proof: the real case. First, remark that since the entries of U_N are in $\{0,1\}$, then for any *-test graph T in one variable, $\tau_N^0\big[T(U_N)\big] = \tau_N^0\big[\tilde T(U_N)\big]$ where $\tilde T$ is obtained by

- reversing the orientation of edges labelled x^* and replacing this label by x,
- reducing positive multiplicity of oriented edges to one.

Hence, we can only consider all test graphs in one variable whose multiplicity of oriented edges is one.

Moreover, each row and column of U_N has a single nonzero entry. Hence, $\tau_N^0\big[T(U_N)\big]$ is zero as soon as two distinct edges leave (or start from) a same vertex. Hence, there are only two kinds of test graphs that possibly contribute: for any $K \geqslant 1$,

- the test graph T_K^c with vertices $1, \dots, K$ and edges $(1,2), \dots, (K-1,K), (K,1)$ (c stands for closed path).
- the test graph T_K^o with vertices $1, \dots, K$ and edges $(1,2), \dots, (K-1,K)$ (o stands for open path).

Let σ_N be the random permutation associated to U_N. Then, $\mathbb{E}\big[\tau_N^0\big[T_K^c(U_N)\big]\big]$ is the probability that a given integer i in $\{1, \dots, N\}$ belongs to a cycle of σ_N of length K. By a straightforward computation, this probability is

$$\frac{N-1}{N} \times \frac{N-2}{N-1} \times \dots \times \frac{N-K}{N-K-1} \times \frac{1}{N},$$

which is of order $\frac{1}{N}$. Then we get

$$\mathbb{E}\Big[\tau_N^0\big[T_K^c(U_N)\big]\Big] \underset{N\to\infty}{\longrightarrow} 0.$$

At the contrary, $\mathbb{E}\big[\tau_N^0\big[T_K^o(U_N)\big]\big]$ is the probability that a given integer i in $\{1,\dots,N\}$ belongs to a cycle of σ_N of length bigger than K. By the above, one has

$$\mathbb{E}\Big[\tau_N^0\big[T_K^o(U_N)\big]\Big] \underset{N\to\infty}{\longrightarrow} 1.$$

Generalities about Hadamard product of matrices invariant by permutation
Consider a random matrix H_N written $H_N = V_N \circ M_N$, where V_N and M_N are independent random matrix, M_N is invariant in law by permutation and $\circ$ denotes the Hadamard (entry-wise) product. Then, for any *-test graph $T=(V,E,\varepsilon)$ in one variable, one has

$$\begin{aligned}
\tau_N^0\big[T(H_N)\big] &= \frac{1}{N}\sum_{\substack{\phi:V\to\{1,\dots,N\}\\ \text{injective}}} \mathbb{E}\Big[\prod_{e\in E} H_N^{\varepsilon(e)}\big(\phi(e)\big)\Big]\\
&= \frac{1}{N}\sum_{\substack{\phi:V\to\{1,\dots,N\}\\ \text{injective}}} \mathbb{E}\Big[\prod_{e\in E} V_N^{\varepsilon(e)}\big(\phi(e)\big)\Big]\times \mathbb{E}\Big[\prod_{e\in E} M_N^{\varepsilon(e)}\big(\phi(e)\big)\Big]\\
&= \mathbb{E}\Big[\tau_N^0\big[T(V_N)\big]\Big]\times \delta_N^0\big[T(M_N)\big],
\end{aligned}$$

where $\delta_N^0\big[T(M_N)\big] = \mathbb{E}\Big[\prod_{e\in E} M_N^{\varepsilon(e)}\big(\phi(e)\big)\Big]$ which does not depend on the injective map ϕ. If in expectation V_N has a limiting distribution of traffics and M_N has a limiting distribution of graphons (i.e. $\delta_N^0\big[T(M_N)\big]$ converges for any T, see Section 0.3), then H_N has a limiting distribution of traffics.

Proof: the complex case. The matrix U_N can be written as the Hadamard product $V_N\circ M_N$, where V_N is a uniform permutation matrix independent of $M_N = (e^{i\theta_{n,m}})_{n,m=1,\dots,N}$. For any *-test graph $T=(V,E,\varepsilon)$ in one variable, by the observation above

$$\mathbb{E}\Big[\tau_N^0\big[T(U_N)\big]\Big] = \mathbb{E}\Big[\tau_N^0\big[T(V_N)\big]\Big]\times \delta_N^0\big[T(M_N)\big],$$

where $\delta_N^0\big[T(M_N)\big] = \mathbb{E}\Big[\prod_{e\in E} M_N^{\varepsilon(e)}\big(\phi(e)\big)\Big]$ for any ϕ. Since $\mathbb{E}[e^{ki\theta_{n,m}}] = \delta_{k0}$ for any $n,m = 1,\dots,N$ and k in $\mathbb{Z}$, we get

$$\delta_N^0\big[T(M_N)\big] = \mathbf{1}_{T \text{ is Hermitian}},$$

where T is Hermitian means that for a given undirected edge e of T, the number of directed edges corresponding to e in one direction and labelled x is equal to the number of directed edges in the opposite direction and labelled x^*. We get the expected result as a result of the real case. □

4.3 Diagonal distribution of traffics associated to a measure on $\mathbb{C}^p$

The following kind of distributions are those of what we called "traffic-incarnations" of commutative random variables.



Figure 10: Three flowers, i.e. *-test graphs with only one edges

Definition 4.6 (Diagonal distribution of traffics).
Let μ be a probability measure on $\mathbb{C}^p$ having all its moments. Let $\mathbf{Z}=(Z_1,\dots,Z_p)$ be a family of random variables distributed according to μ. We call diagonal distribution of traffics associated to μ is the map

$$\begin{array}{cccc}
\tau_\mu: & \mathcal{G}\langle \mathbf{z},\mathbf{z}^*\rangle & \to & \mathbb{C}\\
 & T=(V,E,\gamma,\varepsilon) & \mapsto & \tau_\mu[T] = \mathbb{E}\Big[\prod_{e\in E} Z_{\gamma(e)}^{\varepsilon(e)}\Big],
\end{array}$$

(we use the notation $Z_k^* = \bar{Z}_k$, $k = 1, \dots, p$*).*

In other words, $\tau_\mu[T]$ can be computed by replacing the variables $\mathbf{z}, \mathbf{z}^*$, by the random variable $\mathbf{Z}, \bar{\mathbf{Z}}$ and taking the expectation of the product of the random variables on the edges of T, without taking account into its geometry. The distribution of traffics τ_μ can be easily expressed in terms of its injective version: for any *-test graph $T = (V, E, \gamma, \varepsilon)$ in p variables,

$$\tau_\mu^0[T] = \begin{cases} \mathbb{E}\Big[\prod_{e\in E} Z_{\gamma(e)}^{\varepsilon(e)}\Big] & \text{if } T \text{ has only one vertex } (T \text{ is a "flower"}), \\ 0 & \text{otherwise.} \end{cases}$$

The knowledge of the diagonal distribution of traffics associated to μ is equivalent to the knowledge of its joint moments, that is the number $\mathbb{E}[Z_1^{n_1}, \dots, Z_p^{n_p} \bar{Z}_1^{m_1}, \dots, \bar{Z}_p^{m_p}]$, $n_1, \dots, n_p, m_1, \dots, m_p \geqslant 0$. Hence, τ_μ characterizes μ at soon as μ is characterized by its moments.

Let D_N be an N by N diagonal matrix with complex diagonal entries $\lambda_1, \dots, \lambda_N$. Then its distribution of traffics is the diagonal distribution of traffics associated to the empirical sample distribution of $(\lambda_1, \dots, \lambda_N)$, that is the probability measure

$$\mathcal{L}_{D_N} = \frac{1}{N} \sum_{i=1}^{N} \delta_{\lambda_i}. \tag{4.6}$$

In particular, D_N has a limiting distribution of traffics on $\mathcal{G}\langle \mathbf{z}, \mathbf{z}^* \rangle$ if and only if the measure $\mathcal{L}_{D_N}$ converges in moments.

Part III

General distributions of traffics

We define a notion of space of traffics. It consists of a *-algebra $(\mathcal{A}, .^*)$ endowed with a formal distribution of traffics τ on *-test graphs labelled by elements in $\mathcal{A}$, with the additional condition that τ satisfies a certain condition of positivity. To state this condition, we need to introduce the non commutative *-algebra $\mathbb{C}\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle$, which plays the role of polynomials in space of traffics.

For the task of this article, the reader which is not familiar with the construction of this Part can skip Sections 5 to 7 and only retains the operations on traffics described in Section 8.

5 A *-algebra of multirooted *-test graphs

Definition 5.1 (multirooted *-test graph).
A multirooted *-test graph is a *-test graph with inputs and outputs in certain of its vertices (in a same vertex, they may be many of them. Formally, it consists in a triplet $t = (T, \mathbf{i}, \mathbf{j})$ *where* T *is a* **-test graph and* $\mathbf{i}$ *and* $\mathbf{j}$ *are tuples of vertices of* T*, respectively referred as the sequence of inputs and of outputs.*

The set of all multirooted *-test graphs is denoted by $\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle$ (this set is up to isomorphisms of graphs that preserve the roots). For any $m, n \geqslant 1$, a multirooted *-test graph with m inputs and n inputs is called an $(m,n)^*$-test graph, and the set of such graphs is denoted by $\mathcal{T}_{m,n}\langle \mathbf{x}, \mathbf{x}^* \rangle$. The set of $(n,n)^*$-test graphs is simply denoted $\mathcal{T}_n\langle \mathbf{x}, \mathbf{x}^* \rangle$.

Definition 5.2 (The *-algebra $\mathbb{C}\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle$).
We set $\mathbb{C}\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle$ *the* $\mathbb{C}$*-vector space of finite linear combination of elements of* $\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle$ *with coefficients in* $\mathbb{C}$*. For any* $m, n \geqslant 1$*, we define* $\mathbb{C}\mathcal{T}_{m,n}\langle \mathbf{x}, \mathbf{x}^* \rangle$ *as the vector space of finite linear combinations of elements of* $\mathcal{T}_{m,n}\langle \mathbf{x}, \mathbf{x}^* \rangle$*, and we denote* $\mathbb{C}\mathcal{T}_n\langle \mathbf{x}, \mathbf{x}^* \rangle = \mathbb{C}\mathcal{T}_{n,n}\langle \mathbf{x}, \mathbf{x}^* \rangle$*. In particular,*

$$\mathbb{C}\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle := \bigoplus_{n,m \geqslant 1} \mathbb{C}\mathcal{T}_{m,n}\langle \mathbf{x}, \mathbf{x}^* \rangle.$$

Operations on $\mathbb{C}\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle$:

- **Involution:** Let $t = (T, \mathbf{i}, \mathbf{j})$ be a $(m,n)^*$-test graph. We define its conjugate transpose as the $(n,m)^*$-test graph $t^* = (T^*, \mathbf{j}, \mathbf{i})$, where T^* is the *-test graph obtained by reversing the direction of the edges of T, and replacing labels $\mathbf{x}$ by $\mathbf{x}^*$ and vice versa. We extend $.^*$ into an antilinear map $\mathbb{C}\mathcal{T}_{m,n}\langle \mathbf{x}, \mathbf{x}^* \rangle \to \mathbb{C}\mathcal{T}_{n,m}\langle \mathbf{x}, \mathbf{x}^* \rangle$ and then, into an antilinear involution on $\mathbb{C}\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle$ and on $\mathbb{C}\mathcal{T}_n\langle \mathbf{x}, \mathbf{x}^* \rangle$ for any $n \geqslant 1$.
- **Product:** Let now be $t = (T, \mathbf{i}, \mathbf{j})$ and $t' = (T', \mathbf{i}', \mathbf{j}')$ be (l,m) and $(m',n)^*$-test graphs respectively. We define the product tt' as 0 if $m \neq m'$, and otherwise as the $(l,n)^*$-test graph $(\tilde{T}, \mathbf{i}, \mathbf{j}')$, where $\tilde{T}$ is the *-test graph obtained by considering two disjoint copies of T and T' and identifying j_k and i'_k, $k = 1, \dots, m$. This define a bilinear map $\mathbb{C}\mathcal{T}_{l,m}\langle \mathbf{x}, \mathbf{x}^* \rangle \to \mathbb{C}\mathcal{T}_{m',n}\langle \mathbf{x}, \mathbf{x}^* \rangle$, and then a distributive product on $\mathbb{C}\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle$ and on $\mathbb{C}\mathcal{T}_n\langle \mathbf{x}, \mathbf{x}^* \rangle$ for any $n \geqslant 1$. See examples in Figure 11.

Endowed with their product and their anti-linear involution, the vector spaces $\mathbb{C}\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle$ and $\mathbb{C}\mathcal{T}_n\langle \mathbf{x}, \mathbf{x}^* \rangle$ are then *-algebras. The algebra $\mathbb{C}\mathcal{T}_1\langle \mathbf{x}, \mathbf{x}^* \rangle$ is unital.

6 Distributions of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$

Let τ be a map $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle \to \mathbb{C}$. We define the linear form

$$\Phi^{(\tau)} : \bigoplus_{n \geqslant 1} \mathbb{C}\mathcal{T}_n\langle \mathbf{x}, \mathbf{x}^* \rangle \to \mathbb{C}$$



Figure 11: Top : a $(1,2)^*$-test graph $t = \big(T, (i), (j_1, j_2)\big)$, a $(2,1)^*$-test graph $t' = \big(T', (i_1, i_2), (j)\big)$ and their product tt'. Input vertices are represented in a white rectangle, output vertices are represented in a black one. Bottom: t^*, t'^* and $t't$.

by: for any $t = (T, \mathbf{i}, \mathbf{j})$ in $\mathcal{G}^{(n)}\langle \mathbf{x}, \mathbf{x}^* \rangle$,

$$\Phi^{(\tau)}(t) = \tau[\tilde{T}], \tag{6.1}$$

where $\tilde{T}$ is the *-test graph obtained from T by identifying the k-th input of t and the k-th output, $k = 1, \dots, n$. We extend $\Phi^{(\tau)}$ on $\mathbb{C}\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle$ by $\Phi^{(\tau)}(t) = 0$ for any t in $\mathbb{C}\mathcal{T}_{m,n}\langle \mathbf{x}, \mathbf{x}^* \rangle$, $m \neq n$.

Definition 6.1 (Distributions of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$)**.**
We say that τ is a distribution of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ whenever $\Phi^{(\tau)}$ is a state on $\mathbb{C}\mathcal{T}\langle \mathbf{x}, \mathbf{x}^* \rangle$, that is

$$\Phi^{(\tau)}(t^* t) \geqslant 0, \quad \forall t. \tag{6.2}$$

Notice that such a state $\Phi^{(\tau)}$ is always tracial.

Proposition 6.2. *A distribution of traffics of a family of matrices is a distribution of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$. Hence, a limiting distribution of traffics of matrices on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ is a distribution of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$.*

Proof. The proof is inspired by the constructions in [20, Theorem 11]. Let $\mathbf{A}_N = (A_1, \dots, A_p)$ be a family of complex matrices. First, by definition of the product of multi-rooted *-test graphs, it is enough to prove that for any t in $\mathbb{C}\mathcal{T}_{m,n}\langle \mathbf{x}, \mathbf{x}^* \rangle$, one has $\Phi^{(\tau_{\mathbf{A}_N})}(t^* t) \geqslant 0$. Consider t in $\mathbb{C}\mathcal{T}_{m,n}\langle \mathbf{x}, \mathbf{x}^* \rangle$ and write

$$t = \sum_{k=1}^{d} \alpha_k t_k,$$

where $\alpha_1, \dots, \alpha_d \in \mathbb{C}$ and for any $k = 1, \dots, d$, $t_k = (T_k, \mathbf{i}_k, \mathbf{j}_k) \in \mathcal{T}_{m,n}\langle \mathbf{x}, \mathbf{x}^* \rangle$. Let k in $\{1, \dots, d\}$ and write $T_k = (V_k, E_k, \gamma_k, \varepsilon_k)$, $\mathbf{i}_k = (i_1^{(k)}, \dots, i_m^{(k)})$ and $\mathbf{j}_k = (j_1^{(k)}, \dots, j_n^{(k)})$. Consider the set $V_k^{(0)}$ of vertices in $\mathbf{i}_k$ or in $\mathbf{j}_k$. We denote by $\xi_1, \dots, \xi_N$ the canonical basis of $\mathbb{R}^N$. Define the operator $L_k^{(N)} : (\mathbb{R}^N)^{\otimes m} \to (\mathbb{R}^N)^{\otimes n}$ by: for any $\phi_0 : V_k^{(0)} \to \{1, \dots, N\}$,

$$\begin{aligned} &\langle \xi_{\phi_0(j_1^{(k)})} \otimes \cdots \otimes \xi_{\phi_0(j_n^{(k)})}, L_k^{(N)} \xi_{\phi_0(i_1^{(k)})} \otimes \cdots \otimes \xi_{\phi_0(i_m^{(k)})} \rangle \\ &= \frac{1}{N} \sum_{\phi : V_k \to \{1, \dots, N\}} \prod_{e \in E} A_{\gamma}(e)^{\varepsilon(e)}\big(\phi(e)\big), \end{aligned}$$

where the sum is over all $\phi : V_k \to \{1, \dots, N\}$ such that $\phi_{|V_k^{(0)}} = \phi_0$, and $\langle \xi_{i_1} \otimes \cdots \otimes \xi_{i_n}, L_k^{(N)} \xi_{i_1} \otimes \cdots \otimes \xi_{i_m} \rangle = 0$ otherwise. Now, denote

$$L_N = \sum_{k=1}^{d} \alpha_k L_k^{(N)}.$$

Then, one has

$$\begin{aligned}
\Phi^{(\tau_{\mathbf{A}_N})}(t^*t) &= \sum_{i,j=1}^{d} \bar\alpha_i \alpha_j \Phi^{(\tau_{\mathbf{A}_N})}(t_i^* t_j) \\
&= \frac{1}{N} \sum_{i,j=1}^{d} \bar\alpha_i \alpha_j (\mathrm{Tr} \otimes \cdots \otimes \mathrm{Tr})\big[(L_i^{(N)})^* L_j^{(N)}\big] \\
&= \frac{1}{N} (\mathrm{Tr} \otimes \cdots \otimes \mathrm{Tr})[L_N^* L_N] \geqslant 0.
\end{aligned}$$

□

Proposition 6.3. *Let μ be a measure on $\mathbb{C}^p$ having all its moments. Then, the diagonal distribution of traffics τ_μ associated to μ is a distribution of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$.*

Proof. The map τ_μ is the distribution of traffics of a family of random variable distributed according to μ, seen as a family of 1 by 1 matrices. □

7 Space of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$

We mimic the observations made for the space of N by N matrices. We define algebras of traffics, and then spaces of traffics (which are a ”completed“ version of algebras of traffics).

Definition 7.1 (Algebra of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$)**.**
An algebra of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ is a triplet $(\mathcal{A}, .^*, \tau)$, where

- *$\mathcal{A}$ is a (non commutative) unital algebra over $\mathbb{C}$,*
- *$.^*$ is an antilinear involution such that $(ab)^* = b^* a^*$ for any a, b in $\mathcal{A}$.*
- *τ is a map, called a trace (of traffics) on $\mathcal{A}$,*

$$\begin{array}{rccc} \tau : & \cup_{p\geqslant 1}\big(\mathcal{A}^p \times \mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle\big) & \to & \mathbb{C} \\ & (\mathbf{a}, T) & \mapsto & \tau\big[T(\mathbf{a}, \mathbf{a}^*)\big] := \tau_{\mathbf{a}}[T], \end{array} \tag{7.1}$$

where $\tau_{\mathbf{a}} : \mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle \to \mathbb{C}$ is a distribution of traffics (called the distribution of traffics of **a***), such that τ satisfies the following properties: $p \geqslant 1$ is an integer, $a_1, \dots, a_p, a, b$ are in $\mathcal{A}$, $T = (V, E, \gamma)$ is in $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$.*

1. **Permutation of the indices:** *for any permutation π of $\{1, \dots, p\}$,*

$$\tau\big[T(a_{\pi(1)}, \dots, a_{\pi(p)})\big] = \tau\big[\tilde T(\mathbf{a})\big], \tag{7.2}$$

   where $\tilde T = (V, E, \pi \circ \gamma)$.

2. **Role of the unit:**

$$\tau\big[T(\mathbf{1}, a_1, \dots, a_p)\big] = \tau\big[\tilde T(a_1, \dots, a_p)\big], \tag{7.3}$$

   where $\tilde T$ is the *-test graph obtained by merging adjacent vertices of T joined by an edge labelled **1**.

3. **Role of the involution:**

$$\tau\big[T(a^*, a_1, \dots, a_p)\big] = \tau\big[\tilde T(a, a_1, \dots, a_p)\big], \tag{7.4}$$

   where $\tilde T$ is the *-test graph obtained replace the label of the first variable by its adjoint and vice versa.

4. **Compatibility with the linearity:** *for any complex number* λ,

$$\tau\big[T(\lambda a_1, a_2, \dots, a_p)\big] = \lambda^{\text{Card } \{\gamma^{-1}(1)\}}\tau\big[T(\mathbf{a})\big]. \tag{7.5}$$

and

$$\tau\big[T(a_1, a_2, \dots, a_p, a+b)\big] = \sum_{\tilde T} \tau\big[\tilde T(a_1, , \dots, a_p, a, b)\big], \tag{7.6}$$

where the sum is over **-test graphs* $\tilde T$ *obtained from* T *by choosing for each edge labeled* $(a+b)^\varepsilon$ *a new label among* a^ε *and* b^ε, $\varepsilon \in \{1, *\}$.

5. **Compatibility with the product:**

$$\tau\big[T(a_1, a_2, \dots, a_p, ab)\big] = \tau\big[\tilde T(a_1, , \dots, a_p, a, b)\big], \tag{7.7}$$

where $\tilde T = (\tilde V, \tilde E, \tilde\gamma)$ *is obtained by replacing each edge* $e = (v_1, v_2)$ *such that* $\gamma(e) = p$ *by a path* $e_1 \circ e_2$, *where* $e_1 = (v_1, v_e)$ *and* $e_2 = (v_e, e_2)$, v_e *being a new vertex, linked only at* v_1 *and* v_2 *by* e_1 *and* e_2 *respectively. We set* $\tilde\gamma(e_1) = p$ *and* $\tilde\gamma(e_2) = p+1$.

6. **Consistency:** *if for any edge* e *in* E, $\gamma(e) \neq p$, *then one has*

$$\tau\big[T(\mathbf{a})\big] = \tau\big[\tilde T(a_1, , \dots, a_{p-1})\big],$$

where $\tilde T = (V, E, \tilde\gamma)$, *and* $\tilde\gamma$ *is as* γ *but with range reduced to* $\{1, \dots, p-1\}$.

Let $(\mathcal{A}, .^*, \tau)$ be an algebra of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle$. We define for any a in $\mathcal{A}$,

$$\Phi(a) = \tau\big[T(a)\big], \tag{7.8}$$

where T is the *-test graph with one vertex and one edge (the leftmost test graph in Figure 5). By Definition 6.1 and Formulae (7.7) and (7.4), for any a in $\mathcal{A}$,

$$\Phi(aa^*) \geqslant 0. \tag{7.9}$$

In other word, $\mathcal{A}$ is endowed with a state, and hence is a *-probability space.

Let $(\mathcal{A}, .^*, \tau)$ be an algebra of traffics. Consider the *-algebra $\mathbb{C}\mathcal{T}_1\langle \mathcal{A}\rangle$, formally defined as $\mathbb{C}\mathcal{T}_1\langle \mathbf{x}, \mathbf{x}^*\rangle$, but with labels in $\mathcal{A}$. This algebra is an algebra of traffics when equipped with τ. We identify an element a in $\mathcal{A}$ with $t_a = \big(T, (i), (j)\big)$, where T is the graph with two vertices i and j and one edge from i to j labelled a (see Figure 12, up-leftmost picture). It turns out that the map $a \mapsto t_a$ is an injective morphism of algebra preserving τ, and hence one can see $\mathcal{A}$ as a sub-algebra of traffics of $\mathbb{C}\mathcal{T}_1\langle \mathcal{A}\rangle$. Quotient the algebra $\mathbb{C}\mathcal{T}_1\langle \mathcal{A}\rangle$ by elements a such that $\Phi^{(\tau)}(aa^*) = 0$. With a small abuse of notation, still denote $\mathbb{C}\mathcal{T}_1\langle \mathcal{A}\rangle$ this algebra.

Definition 7.2 (Space of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle$)**.**
A space of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle$ *is an algebra of traffics on* $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle$ *such that the homomorphism* $a \mapsto t_a$ *described above is an isomorphism* $\mathcal{A} \to \mathbb{C}\mathcal{T}_1\langle \mathcal{A}\rangle$. *Elements of a space of traffics on* $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle$ *are called traffics.*

Definition 7.3 (Convergence in distribution of traffics)**.**
Let $\mathbf{t}, \mathbf{t}_n, n \geqslant 1$ *be families of traffics of the same cardinal in a space of traffics* $(\mathcal{A}, .^*, \tau)$. *Let* $\mathcal{D}$ *be a domain*

$$\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^*\rangle \subset \mathcal{D} \subset \mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle.$$

We say that $\mathbf{t}_n$ *converges to* $\mathbf{t}$ *in distribution of traffics on* $\mathcal{D}$ *whenever* $\tau_{\mathbf{t}_n}$ *converges pointwise on* $\mathcal{D}$ *to* $\tau_{\mathbf{t}}$, *i.e.*

$$\forall T \in \mathcal{D}, \quad \tau[T(\mathbf{t})] = \lim_{n\to\infty} \tau\big[T(\mathbf{t}_n)\big] \quad \textit{exists.} \tag{7.10}$$



Figure 12: Some traffics in $\mathbb{C}\mathcal{T}_1\langle\mathcal{A}\rangle$. Black edges are labelled a in $\mathcal{A}$, red ones are labelled b in $\mathcal{A}$. Top, from left to right: a, a^*, a^t and $\bar{a}$. Bottom, from left to right: ab, $a\circ b$ and Δ_a.

8 Remarkable operations on traffics

All the definitions of this Sections are summed up in Figure 12.

The Hadamard product $a\circ b$:
Let a, b be two traffics in a space of traffics $(\mathcal{A}, .^*, \tau)$ on $\mathcal{G}\langle\mathbf{x},\mathbf{x}^*\rangle$. The Hadamard product of a and b, denoted $a\circ b$, is the traffic corresponding to $L=(T,s,t)$ in $\mathbb{C}\mathcal{T}_1\langle\mathcal{A}\rangle$, where T is the test graph with two vertices s and t and two edges from s to t, one labelled a and the other one labelled b.

The Hadamard product is symmetric. The Hadamard product of matrices seen as traffics is well their entry-wise product.

The projection on the diagonal:
Let a be a traffic in a space of traffics $(\mathcal{A}, .^*, \tau)$ on $\mathcal{G}\langle\mathbf{x},\mathbf{x}^*\rangle$. The projection of a on the diagonal is the traffic $\Delta_a = a\circ\mathbf{1}$, where $\mathbf{1}$ is the unit of $\mathcal{A}$. Equivalently, Δ_a is the traffic corresponding to $L=(T,s,s)$ in $\mathbb{C}\mathcal{T}_1\langle\mathcal{A}\rangle$, where T is the test graph with one vertex s and one edge labelled a.

The projection of a matrix A_N on the diagonal is the diagonal matrix Δ_{A_N} of its diagonal elements. The sub-algebra of $\mathcal{A}$ spanned by projections on the diagonal of elements of $\mathcal{A}$ is a commutative sub-algebra of $\mathcal{A}$. For any a in $\mathcal{A}$ and any $k,l\geqslant 1$, one has

$$\Phi\big[\Delta_a^k\Delta_a^{*l}\big] = \Phi\big[\ \underbrace{a\circ\cdots\circ a}_{k\text{ times}}\circ\underbrace{a^*\circ\cdots\circ a^*}_{l\text{ times}}\ \big].$$

Transpose and conjugate:
Let a be a traffic in a space of traffics $(\mathcal{A}, .^*, \tau)$ on $\mathcal{G}\langle\mathbf{x},\mathbf{x}^*\rangle$. The transpose of a, denoted a^t, is the traffic corresponding to $L=(T,s,t)$ in $\mathbb{C}\mathcal{T}_1\langle\mathcal{A}\rangle$, where T is the test graph with two vertices s and t and one edge from t to s labelled a. The conjugate of a is the traffic $\bar{a}=(a^*)^t$.

For matrices, transpose and conjugate are the classical operations.

9 Space of traffics on $\mathcal{G}_{cyc}\langle\mathbf{x},\mathbf{x}^*\rangle$

It will be convenient to have a cyclic version of space of traffics. We set $\mathcal{T}_{1,cyc}\langle\mathbf{x},\mathbf{x}^*\rangle$ as the set of all $t=\big(T,(i),(j)\big)$ in $\mathcal{T}_1\langle\mathbf{x},\mathbf{x}^*\rangle$ such that there exists a path from i to j in T visiting each edges once in the sense of their orientation. We denote by $\mathbb{C}\mathcal{T}_{1,cyc}\langle\mathbf{x},\mathbf{x}^*\rangle$ the vector space of finite linear combination of elements of $\mathcal{T}_{1,cyc}\langle\mathbf{x},\mathbf{x}^*\rangle$ with coefficients in $\mathbb{C}$

Let $\tau:\mathcal{G}_{cyc}\langle\mathbf{x},\mathbf{x}^*\rangle\to\mathbb{C}$ be a formal distribution of traffics. Define the linear from $\Phi^{(\tau)}$ on $\mathbb{C}\mathcal{T}_{1,cyc}\langle\mathbf{x},\mathbf{x}^*\rangle$ as in (6.1). We say that τ is a distribution of traffics on $\mathcal{G}_{cyc}\langle\mathbf{x},\mathbf{x}^*\rangle$ whenever $\Phi^{(\tau)}$ is a state on $\mathcal{Q}\mathcal{G}^{(1)}_{cyc}\langle\mathbf{x},\mathbf{x}^*\rangle$. A space of traffics on $\mathcal{G}_{cyc}\langle\mathbf{x},\mathbf{x}^*\rangle$ is define as a space of traffics on $\mathcal{G}\langle\mathbf{x},\mathbf{x}^*\rangle$ where we have replaced $\mathcal{G}\langle\mathbf{x},\mathbf{x}^*\rangle$ by $\mathcal{G}_{cyc}\langle\mathbf{x},\mathbf{x}^*\rangle$ and $\mathcal{T}_1\langle\mathbf{x},\mathbf{x}^*\rangle$ by $\mathcal{T}_{1,cyc}\langle\mathbf{x},\mathbf{x}^*\rangle$.

10 A domain of sequential compactness by Mingo and Speicher

Motivated by the asymptotic properties of product of large matrices, Mingo and Speicher [20] have introduced the notion of graph of matrices and the sum associated to such objects. In our language of traffics, a graph of matrices is a test graph whose variables have been replaced by matrices. When connected, it is also equivalent to an element of the algebra $\mathbb{C}\mathcal{T}_1\langle \mathrm{M}_N(\mathbb{C})\rangle$ where inputs and outputs has been merged. The sum associated to a graph of matrices in [20] is then the trace of this test graph in the considered matrices.

In the context of distribution of traffics, their main result solves the following question: finding a maximal domain $\mathcal{D}$ for which any family of matrices bounded in operator norm converges in distribution of traffics up to a subsequence.

10.1 Two edge connected and proper test graphs

We follow the definitions in [20].

Definition 10.1 (Cutting edge and two-edge connected component of a graph)**.**

1. *A cutting edge of a graph is an edge whose removal would result into disconnected subgraphs.*
2. *A two-edge connected graph is a graph without cutting edges.*
3. *A two-edge connected component of a graph is a subgraph which is two-edge connected and cannot be enlarged to a bigger two-edge connected subgraph.*

Let G a graph. Its forest of two-edge connected components $\mathfrak{F}(G)$ is defined as follow (see Figure 13): the vertices of $\mathfrak{F}(G)$ consists in the two-edge connected components of G and two distinct vertices of $\mathfrak{F}(G)$ are connected by an edge if there is a cutting edge between vertices from the two corresponding components in G. Hence $\mathfrak{F}(G)$ is always a forest, i.e. a graph without cycles, and if G is connected then $\mathfrak{F}(G)$ is actually a tree. For a *-graph in several variables $T = (G, \gamma, \varepsilon)$, we define $\mathfrak{F}(T) = \mathfrak{F}(G)$.



Figure 13: Left: a graph G whose two-edge connected components has been encircled. Right: its forest $\mathfrak{F}(G)$ of two-edge connected components.

Definition 10.2 (Proper test graphs)**.**

1. *A tree is trivial if it consists in only one vertex. A leaf of a non-trivial tree is a vertex which meets only one edge. By convention, we say that the trivial tree has two leafs.*
2. *A proper test a graph is a test graph whose forest of two-edge connected components has two leaves (see Figure 14). The set of all proper test graphs in p variables is denote by $\mathcal{G}_{pr}\langle x_1, \dots, x_p\rangle$.*

As before we use the shortcuts $\mathcal{G}_{pr}\langle \mathbf{x}\rangle$ and $\mathcal{G}_{pr}\langle \mathbf{x}, \mathbf{x}^*\rangle$. In Figure 14 is drawn a proper test graph. In Figure 5, all but the rightmost test graph are proper. Notice that we have the inclusions

$$\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^*\rangle \subset \mathcal{G}_{pr}\langle \mathbf{x}, \mathbf{x}^*\rangle \subset \mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle. \tag{10.1}$$



Figure 14: A proper test graph. Its two-edge connected components have been encircled.

10.2 A domain of sequential compactness of the unit ball

Theorem 10.3 (Sharp bounds for the trace of test graphs in matrices, [20])**.**
Let T be a *-test graph in p variables. Let $\mathfrak{F}(T)$ be its forest of two-edge connected components and denote by $\mathfrak{r}(T)$ its number of leafs. Then, for any family $\mathbf{A}_N = (A_1^{(N)}, \dots, A_p^{(N)})$ of N by N matrices,

$$\Big| \tau_N\big[T(\mathbf{A}_N)\big] \Big| \leqslant N^{\frac{\mathfrak{r}(T)}{2}-1} \prod_{e \in E} \| A_{\gamma(e)}^{(N)} \|, \tag{10.2}$$

where $\| \cdot \|$ stands for the operator norm.

From Theorem 10.3, we deduce immediately the following corollary.

Corollary 10.4 (Sequentially compactness)**.**
Let $\mathbf{A}_N$ be a family of N by N matrices of operator norm uniformly bounded. Then, there exists a subsequence of $(\mathbf{A}_N)_{N \geqslant 1}$ having a limiting distribution of traffics with domain $\mathcal{G}_{pr}\langle \mathbf{x}, \mathbf{x}^* \rangle$.

Contrariwise, as notice in [20, Section 5], if we consider the matrix

$$A_N = \frac{1}{\sqrt{N}} \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & \cdots & 0 \end{pmatrix} \tag{10.3}$$

then $\|A_N\| = 1$ and for any non proper graph T

$$\Big| \tau_N\big[T(A_N)\big] \Big| \underset{N \to \infty}{\longrightarrow} \infty. \tag{10.4}$$

For instance, with the rightmost test graph T of Figure 5, one has

$$\tau_N\big[T(A_N)\big] = \sqrt{N}. \tag{10.5}$$

Part IV

The freeness of traffics

The notion of freeness emerges naturally from the analysis of the distribution of traffics of random matrices (see Theorem 12.1). It consists in a rule that allows to form a joint distribution of families $\mathbf{a}_1$ and $\mathbf{a}_2$ of traffics, in terms of the distribution of $\mathbf{a}_1$ and $\mathbf{a}_2$ respectively. This notion generalizes the notion of independence, when random variables are lifted into diagonal traffics (see Proposition 11.2). It is natural to formulate a central limit theorem for the normalized sum of free traffics: the normal traffics turns out to be the sum of a diagonal gaussian traffic and a semicircular traffic, i.e. the limit in distribution of traffics of a Wigner matrix (see Theorem 13.1), these traffics being free.

11 Definition and first examples

11.1 The free product of *-test graphs

Let $\mathbf{x}_1, \dots, \mathbf{x}_p$ be disjoint families of indeterminates. A *-test graph T is said to be a free product of *-test graphs in $\mathbf{x}_1, \dots, \mathbf{x}_p$ whenever it could be obtained by the following construction (bear in mind that a *-test graph can consists in a single vertex with no edge).

> **Step 1:** Consider *-test graphs $T_{1,0}$ in $\mathcal{G}\langle \mathbf{x}_1, \mathbf{x}_1^* \rangle$, ..., $T_{p,0}$ in $\mathcal{G}\langle \mathbf{x}_1, \mathbf{x}_1^* \rangle$. For any of these *-test graphs, chose a vertex (called its root), and identify the roots of the different *-test graphs to form a single *-test graph $T^{(1)}$.
>
> **Step 2:** Let j be in $\{1, \dots, p\}$. For any vertex v_j of $T_{j,0}$ which is not its root, chose a *-test graph $T_{k,0v_j}$ in $\mathcal{G}\langle \mathbf{x}_k, \mathbf{x}_k^* \rangle$ for any $k \neq j$. For any of these *-test graphs, chose a root and identify the vertex v_j with the root of $T_{k,0v_j}$, $k \neq j$. We get a *-test graph $T^{(2)}$.
>
> **Step 3:** Let j be in $\{1, \dots, p\}$ and $k \neq j$. For any vertex v_j of $T_{j,0}$ which is not its root and any vertex w_k of $T_{k,0v_j}$ which is not its root, chose a *-test graph $T_{\ell,0v_jw_k}$ in $\mathcal{G}\langle \mathbf{x}_\ell, \mathbf{x}_\ell^* \rangle$ for any $\ell \neq k$. For any of these *-test graphs, chose a root and identify the vertex w_k with the root of $T_{\ell,0v_jw_k}$, $\ell \neq k$. We get a *-test graph $T^{(3)}$.
>
> **Further steps:** Iterate this process a finite number of times, and then obtain a *-test graph T

The *-test graphs in $\mathcal{G}\langle \mathbf{x}_j, \mathbf{x}_j^* \rangle$, $j = 1, \dots, p$, involved the construction of T are referred as the components of T. See examples in Figures 3 and 15. A free product of *-test graphs is cyclic if and only if its components are cyclic.

11.2 The freeness of a family of traffics

Definition 11.1 (The freeness of traffics).
Let $(\mathcal{A}, .^*, \tau)$ be a space of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ (respectively $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$) and let $\mathbf{a}_1, \dots, \mathbf{a}_p$ be families of traffics in $\mathcal{A}$. The families $\mathbf{a}_1, \dots, \mathbf{a}_p$ are said to be free on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ (respectively $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$) whenever their joint distribution of traffics $\tau_{\mathbf{a}}$ is given by: for any *-test graph T in $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ (respectively $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$),

- *If T is a free product of *-test graph in $\mathcal{G}\langle \mathbf{x}_j, \mathbf{x}_j^* \rangle$, $j = 1, \dots, p$, then*

  $$\tau^0_{\mathbf{a}}[T] = \prod_{\tilde{T}} \tau^0_{\mathbf{a}}[\tilde{T}],$$

  where the sum is over the components of T.

- *otherwise, $\tau^0_{\mathbf{a}}[T] = 0$.*

Remarks:



Figure 15: Some free products of *-test graphs in x (red) and y (red), from left to right: between flowers, between directed lines, between double trees. The fourth picture is a *-test graph between flowers in the variable a (red) and double trees in the variable b (black), prototype of a cyclic *-test graph that contributes in the central limit theorem (see Section 13).

1. When $\mathbf{a}_1, \dots, \mathbf{a}_K$ are free on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ or $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$, their joint distribution of traffics is completely characterized by the marginal distributions $\tau_{\mathbf{a}_1}, \dots, \tau_{\mathbf{a}_K}$.
2. Freeness on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ implies freeness on $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$, the reciprocal is not true (see Proposition 11.3).

11.3 Examples

11.3.1 Independence and diagonal traffics

Proposition 11.2 (Traffic freeness of diagonal traffics is independence)**.**
Let $(x_1, \dots, x_p, y_1, \dots, y_q)$ *be a family of traffics having a diagonal distribution* τ_μ *associated to a probability measure* μ *on* $\mathbb{C}^{p+q}$ *characterized by its moments (see Section 4.3). Then,* $\mathbf{x} = (x_1, \dots, x_p)$ *and* $\mathbf{y} = (y_1, \dots, y_q)$ *are free on* $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ *if and only if* μ *is the law of random variables* $(X_1, \dots, X_p, Y_1, \dots, Y_q)$ *where* $\mathbf{X} = (X_1, \dots, X_p)$ *and* $\mathbf{Y} = (Y_1, \dots, Y_q)$ *are independent.*

Proof. Assume that $\mathbf{x}$ and $\mathbf{y}$ are free. Since μ is characterized by its moments, it is enough to prove that for any commutative monic monomials P and Q,

$$\mathbb{E}\big[P(\mathbf{X}, \bar{\mathbf{X}})Q(\mathbf{Y}, \bar{\mathbf{Y}})\big] = \mathbb{E}\big[P(\mathbf{X}, \bar{\mathbf{X}})\big]\mathbb{E}\big[Q(\mathbf{Y}, \bar{\mathbf{Y}})\big].$$

Write $P = x_1^{n_1}\bar{x}_1^{m_1} \dots x_p^{n_p}\bar{x}_p^{m_p}$ and $Q = y_1^{k_1}\bar{y}_1^{l_1} \dots y_p^{k_p}\bar{y}_p^{l_p}$. Then, we have

$$\mathbb{E}\big[P(\mathbf{X}, \bar{\mathbf{X}})Q(\mathbf{Y}, \bar{\mathbf{Y}})\big] = \tau_\mu[T] = \tau_\mu^0[T],$$

where T is the flower with n_i edges labelled x_i, m_i edges labelled x_i^*, $i = 1, \dots, p$, and k_i edges labelled y_i, l_i edges labelled y_i^*, $i = 1, \dots, q$. A flower is always a free product of flowers in marginal variables, and the freeness relation gives

$$\tau_\mu^0[T] = \tau_\mu^0[T_1]\tau_\mu^0[T_2],$$

where T_1 is the flower with n_i edges labelled x_i, m_i edges labelled x_i^*, $i = 1, \dots, p$, and T_2 is the flower with k_i edges labelled y_i, l_i edges labelled y_i^*, $i = 1, \dots, q$. Since

$$\tau_\mu^0[T_1]\tau_\mu^0[T_2] = \tau_\mu[T_1]\tau_\mu[T_2] = \mathbb{E}\big[P(\mathbf{X}, \bar{\mathbf{X}})\big]\mathbb{E}\big[Q(\mathbf{Y}, \bar{\mathbf{Y}})\big],$$

we get that $\mathbf{X}$ and $\mathbf{Y}$ are independent.

Reciprocally, assume that $\mathbf{X}$ and $\mathbf{Y}$ are independent. The distribution τ_μ is supported on flowers, which are free products of *-test graphs. With the same notation as above, we have to show that $\tau_\mu^0[T] = \tau_\mu^0[T_1]\tau_\mu^0[T_2]$, which is obtained from the independence of $\mathbf{X}$ and $\mathbf{Y}$ observed on monic monomials. ☐

11.3.2 Asymptotic freeness on $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$ of a complex Wigner matrix and its transpose

Proposition 11.3. *Let x be a traffic in a space $(\mathcal{A}, .^*, \tau)$, limit in distribution of traffics of a complex Wigner matrix (see Proposition 4.2). Then, x and x^t are free on $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^* \rangle$ but not in $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$.*



Figure 16: Two test graphs that contribute, at the level of the injective trace, in the limiting distribution of traffics of (X_N, X_N^t). The leftmost test graph is cyclic and is a free product of black and red double trees. The rightmost is not cyclic and is not a free product.

Proof. To prove that x and x^t are not free on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$, consider the test graph T with two vertices i and j and two edges from i to j, one labelled x_1 and the other labelled x_2. Since T is not a free product of test graphs in x_1 and x_2, we get the expected result if we show that $\tau^0[T(x, x^t)] \neq 0$.

Let T' be the test graph analogue of T but where the two edges have different directions. By definition of the transpose, one has

$$\tau\big[T(x, x^t)\big] = \tau\big[T'(x, x)\big] = 1.$$

Let $\tilde{T}$ be the test graph obtained from T by identifying its two vertices. By definition of the injective trace, one has

$$\tau\big[T(x, x^t)\big] = \tau^0\big[T(x, x^t)\big] + \tau^0\big[\tilde{T}(x, x^t)\big].$$

But

$$\tau^0\big[\tilde{T}(x, x^t)\big] = \tau^0\big[\tilde{T}(x, x)\big] = 0,$$

since $\tilde{T}$ is not a double tree, and so we get $\tau^0\big[T(x, x^t)\big] = 1 \neq 0$ Hence, x and x^t are not free in $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$.

Let now T be a cyclic test graph in two variables x_1 and x_2. Let T' be the test graph in one variable obtained from T by reversing the orientation of edges labelled x_2, and relabeling them x_1. Then

$$\tau^0\big[T(x, x^t)\big] = \tau^0\big[T'(x)\big].$$

Notice that a double tree which is cyclic has necessarily twin edges of opposite directions. Hence, we get

- $\tau^0\big[T(x, x^t)\big] = 1$ if T is a double tree whose twin edges have the same label,

- $\tau^0\big[T(x,x^t)\big]=0$ otherwise.

Since x^t has the same distribution as x, we get that x and x^t are free on $\mathcal{G}_{cyc}\langle \mathbf{x},\mathbf{x}^*\rangle$. ☐

Proposition 11.4. *Let u be a traffic in a space $(\mathcal{A},.^*,\tau)$, limit in distribution of traffics of a complex permutation matrix (see Proposition 4.5). Then, u and u^t are not free on $\mathcal{G}_{cyc}\langle \mathbf{x},\mathbf{x}^*\rangle$.*

Proof. Let T be the *-test graph (see Figure 17) with two vertices i and j and

- one edge from i to j labelled x_1,
- one edge from j to i labelled x_1^*,
- one edge from i to j labelled x_2^*,
- one edge from j to i labelled x_2.

We have $\tau^0\big[T(u,u^t)\big]=1$ although T is not a free product of *-test graphs in x_1 and x_2. ☐



Figure 17: The *-test graph T of the proof of Proposition 11.4.

12 An asymptotic freeness theorem for random matrices

12.1 Statement and proof

Theorem 12.1 (The asymptotic freeness of $\mathbf{X}_1^{(N)},\dots,\mathbf{X}_p^{(N)}$ on $\mathcal{G}\langle \mathbf{x},\mathbf{x}^*\rangle$**).**
Let $\mathbf{X}_1^{(N)},\dots,\mathbf{X}_p^{(N)}$ be families of N by N random matrices. Let $\mathbf{x}_1,\dots,\mathbf{x}_p$ be families of traffics in a space of traffics $(\mathcal{A},.^*,\tau)$ on $\mathcal{G}\langle \mathbf{x},\mathbf{x}^*\rangle$. Assume the following.

1. **Statistical independence:**
   The families $\mathbf{X}_1^{(N)},\dots,\mathbf{X}_p^{(N)}$ are statistically independent.
2. **Joint invariance by permutation:**
   For any permutation matrix U_N, and any $j=1,\dots,p$,
$$U_N\mathbf{X}_j^{(N)}U_N^* \overset{\mathcal{L}}{=} \mathbf{X}_j^{(N)}. \tag{12.1}$$
3. **Freeness of $\mathbf{x}_1,\dots,\mathbf{x}_p$:**
   The families of traffics $\mathbf{x}_1,\dots,\mathbf{x}_p$ are free on $\mathcal{G}\langle \mathbf{x},\mathbf{x}^*\rangle$.
4. **Convergence in distribution of traffics on $\mathcal{G}\langle \mathbf{x},\mathbf{x}^*\rangle$:**
   For any $j=1,\dots,p$, in expectation $\mathbf{X}_j^{(N)}$ converges in distribution of traffics to $\mathbf{x}_j$ on $\mathcal{G}\langle \mathbf{x},\mathbf{x}^*\rangle$ and moreover: for any *-test graphs $T_1,\dots,T_n$,
$$\mathbb{E}\Big[\tau_N\big[T_1(\mathbf{X}_j^{(N)})\big]\dots\tau_N\big[T_n(\mathbf{X}_j^{(N)})\big]\Big] \underset{N\to\infty}{\longrightarrow} \tau\big[T_1(\mathbf{x}_j)\big]\dots\tau\big[T_n(\mathbf{x}_j)\big]. \tag{12.2}$$

Then, in expectation, the joint family $(\mathbf{X}_1^{(N)},\dots,\mathbf{X}_p^{(N)})$ converges in distribution of traffics to $(\mathbf{x}_1,\dots,\mathbf{x}_p)$ on $\mathcal{G}\langle \mathbf{x},\mathbf{x}^*\rangle$, and moreover for any *-test graphs $T_1,\dots,T_n$

$$\begin{aligned}&\mathbb{E}\Big[\tau_N\big[T_1(\mathbf{X}_1^{(N)},\dots,\mathbf{X}_p^{(N)})\big]\dots\tau_N\big[T_n(\mathbf{X}_1^{(N)},\dots,\mathbf{X}_p^{(N)})\big]\Big]\\ &\underset{N\to\infty}{\longrightarrow} \tau\big[T_1(\mathbf{x}_1,\dots,\mathbf{x}_p)\big]\dots\tau\big[T_n(\mathbf{x}_1,\dots,\mathbf{x}_p)\big].\end{aligned} \tag{12.3}$$

Proof of Theorem 12.1. By Lemma 3.2, relating the injective and the non injective traces, it is equivalent to prove the convergence of injective traces in (12.3). For clarity of the presentation, in the first three steps of this proof, we prove (12.3) for $n=1$ and for two families $\mathbf{X}_1^{(N)}$ and $\mathbf{X}_2^{(N)}$. The case $n \geqslant 1$, with two families of matrices, is obtain in the fourth step. The general case is obtained by recurrence on the number of families.

Step 1: Splitting the contribution due to $\mathbf{X}_1^{(N)}$ and $\mathbf{X}_2^{(N)}$

Lemma 12.2 (A probabilistic interpretation of the injective trace)**.**

1. *Let V be a finite set. Let $\Phi_N, \tilde{\Phi}_N$ be two independent random maps, uniformly distributed on the set of injective maps $V \to \{1,\dots,N\}$. Let σ_N be a random permutation on $\{1,\dots,N\}$ uniformly distributed and independent of $(\Phi_N, \tilde{\Phi}_N)$. Let ϕ be an deterministic injective map $V \to \{1,\dots,N\}$. Then,*

$$\sigma_N \circ \phi \overset{\mathcal{L}}{=} \Phi_N \quad \textit{and} \quad (\Phi_N, \sigma_N \circ \Phi_N) \overset{\mathcal{L}}{=} (\Phi_N, \tilde{\Phi}_N).$$

2. *Let $T=(V,E,\gamma,\varepsilon)$ be a finite *-graph in p variables and Φ_N as above. Then, for any family $\mathbf{A}_N=(A_1,\dots,A_p)$ of deterministic complex matrices,*

$$\tau_N^0\big[T(\mathbf{A}_N)\big] = \frac{(N-1)!}{(N-|V|)!}\mathbb{E}\Big[\prod_{e\in E} A_{\gamma(e)}^{\varepsilon(e)}\big(\Phi_N(e)\big)\Big].$$

Proof of Lemma 12.2. Claim 1. A direct computation gives: for any $\phi, \tilde{\phi}$ injective maps $V \to \{1,\dots,N\}$,

$$\mathbb{P}\Big(\sigma_N \circ \phi = \tilde{\phi}\Big) = \frac{(N-|V|)!}{N!} = \mathbb{P}\Big(\Phi_N = \tilde{\phi}\Big)$$

and

$$\begin{aligned}\mathbb{P}\Big(\Phi_N = \phi, \sigma_N \circ \Phi_N = \tilde{\phi}\Big) &= \mathbb{P}\Big(\Phi_N = \phi, \sigma_N \circ \phi = \tilde{\phi}\Big) = \mathbb{P}\Big(\Phi_N = \phi\Big)\mathbb{P}\Big(\sigma_N \circ \phi = \tilde{\phi}\Big)\\ &= \mathbb{P}\Big(\Phi_N = \phi\Big)\mathbb{P}\Big(\tilde{\Phi}_N = \tilde{\phi}\Big)\end{aligned}$$

Claim 2. One has

$$\begin{aligned}\tau_N^0\big[T(\mathbf{A}_N)\big] &= \frac{1}{N}\sum_{\substack{\phi: V\to\{1,\dots,N\}\\ \text{injective}}}\prod_{e\in E} A_{\gamma(e)}^{\varepsilon(e)}\big(\phi(e)\big)\\ &= \frac{(N-1)!}{(N-|V|)!}\frac{1}{\text{Card}\left\{\substack{\phi:V\to\{1,\dots,N\}\\ \text{injective}}\right\}}\sum_{\substack{\phi: V\to\{1,\dots,N\}\\ \text{injective}}}\prod_{e\in E} A_{\gamma(e)}^{\varepsilon(e)}\big(\phi(e)\big)\\ &= \frac{(N-1)!}{(N-|V|)!}\mathbb{E}\Big[\prod_{e\in E} A_{\gamma(e)}^{\varepsilon(e)}\big(\Phi_N(e)\big)\Big].\end{aligned}$$

□

Proposition 12.3 (The permutation invariance property)**.** *Let $\mathbf{X}_1^{(N)}$ and $\mathbf{X}_2^{(N)}$ be two families of N by N permutation invariant, independent random matrices: for any $i=1,2$ and any permutation matrix U_N,*

$$\mathbf{X}_i \overset{\mathcal{L}}{=} U_N \mathbf{X}_i U_N^*$$

Let $T=(V,E,\gamma,\varepsilon)$ be a finite *-graph. For $i=1,2$, we denote by $T_i=(V_i,E_i,\gamma_i,\varepsilon_i)$ the *-graph obtained from T by considering only the edges with a label corresponding to the family $\mathbf{X}_i^{(N)}$ and by deleting the vertices that are not attached to any edge after this process. We have

$$\begin{aligned}&\mathbb{E}\Big[\tau_N^0\big[T(\mathbf{X}_1^{(N)},\mathbf{X}_2^{(N)})\big]\Big]\\ &\quad= \frac{(N-|V_1|)!(N-|V_2|)!}{(N-|V|)!(N-1)!}\mathbb{E}\Big[\tau_N^0\big[T_1(\mathbf{X}_1^{(N)})\big]\Big]\mathbb{E}\Big[\tau_N^0\big[T_2(\mathbf{X}_2^{(N)})\big]\Big]. \qquad (12.4)\end{aligned}$$

Proof of Proposition 12.3. Let Φ_N be a uniform random injection $V \to \{1, \dots, N\}$, independent of $(\mathbf{X}_1^{(N)}, \mathbf{X}_2^{(N)})$. Then, by the second item of Lemma 12.2,

$$\mathbb{E}\Big[\tau_N^0\big[T(\mathbf{X}_1^{(N)}, \mathbf{X}_2^{(N)})\big]\Big] = \frac{(N-1)!}{(N-|V|)!}\mathbb{E}\Big[\prod_{e\in E} X_{\gamma(e)}^{\varepsilon(e)}\big(\Phi_N(e)\big)\Big].$$

Let E_1 be the subset of E formed by edges with a label corresponding to the family $\mathbf{X}_1^{(N)}$, and let $E_2 = E \setminus E_1$. Then, one has

$$\mathbb{E}\Big[\prod_{e\in E} X_{\gamma(e)}^{\varepsilon(e)}\big(\Phi_N(e)\big)\Big] = \mathbb{E}\Big[\prod_{e\in E_1} X_{\gamma(e)}^{\varepsilon(e)}\big(\Phi_N(e)\big) \times \prod_{e\in E_2} X_{\gamma(e)}^{\varepsilon(e)}\big(\Phi_N(e)\big)\Big].$$

Let $\tilde\Phi_N$ be distributed as Φ_N and let σ_N be a uniform permutation on $\{1, \dots, N\}$, such that $(\mathbf{X}_1^{(N)}, \mathbf{X}_2^{(N)})$, Φ_N, $\tilde\Phi_N$ and σ_N are independent. Let U_N be the permutation matrix associated to σ_N. By invariance of $\mathbf{X}_2^{(N)}$ by permutation, $U_N\mathbf{X}_2^{(N)}U_N^*$ and $\mathbf{X}_2^{(N)}$ have the same distribution. Since the entry (i,j) of $U_N M_N U_N^*$ is $M_N\big(\sigma_N(i), \sigma_N(j)\big)$, we get by the first item of Lemma 12.2

$$\begin{aligned}\mathbb{E}\Big[\prod_{e\in E} X_{\gamma(e)}^{\varepsilon(e)}\big(\Phi_N(e)\big)\Big] &= \mathbb{E}\Big[\prod_{e\in E_1} X_{\gamma(e)}^{\varepsilon(e)}\big(\Phi_N(e)\big) \times \prod_{e\in E_2} X_{\gamma(e)}^{\varepsilon(e)}\big(\sigma_N\circ\Phi_N(e)\big)\Big]\\ &= \mathbb{E}\Big[\prod_{e\in E_1} X_{\gamma(e)}^{\varepsilon(e)}\big(\Phi_N(e)\big)\Big] \times \mathbb{E}\Big[\prod_{e\in E_2} X_{\gamma(e)}^{\varepsilon(e)}\big(\tilde\Phi_N(e)\big)\Big]\end{aligned}$$

Again, by the second item of Lemma 12.2, one has

$$\begin{aligned}&\mathbb{E}\Big[\prod_{e\in E_1} X_{\gamma(e)}^{\varepsilon(e)}\big(\Phi_N(e)\big)\Big] \times \mathbb{E}\Big[\prod_{e\in E_2} X_{\gamma(e)}^{\varepsilon(e)}\big(\tilde\Phi_N(e)\big)\Big]\\ &= \frac{(N-|V_1|)!(N-|V_2|)!}{(N-1)!^2}\,\mathbb{E}\Big[\tau_N^0\big[T_1(\mathbf{X}_1^{(N)})\big]\Big]\mathbb{E}\Big[\tau_N^0\big[T_2(\mathbf{X}_2^{(N)})\big]\Big].\end{aligned}$$

Hence, we get

$$\begin{aligned}&\mathbb{E}\Big[\tau_N^0\big[T(\mathbf{X}_1^{(N)}, \mathbf{X}_2^{(N)})\big]\Big]\\ &= \frac{(N-|V_1|)!(N-|V_2|)!}{(N-|V|)!(N-1)!}\mathbb{E}\Big[\tau_N^0\big[T_1(\mathbf{X}_1^{(N)})\big]\Big]\mathbb{E}\Big[\tau_N^0\big[T_2(\mathbf{X}_2^{(N)})\big]\Big].\end{aligned}$$

□

In the sequel of this proof, we use the notation of Proposition 12.3.

Step 2: Analysis of the terms due to the random matrices
Fix i in $\{1,2\}$ until the end of this step. By Lemma 3.2, relating the injective and the non injective traces,

$$\tau_N^0\big[T_i(\mathbf{X}_i^{(N)})\big] = \sum_{\pi\in\mathcal{P}(V_i)} \mu_{V_i}(\pi)\ \tau_N\big[\pi(T_i)(\mathbf{X}_i^{(N)})\big]. \tag{12.5}$$

Let $T_{i,k} = (V_{i,k}, E_{i,k}, \gamma_{i,k}, \varepsilon_{i,k}), k = 1, \dots, K_i$, be the connected components of T_i. If π is a partition as in the sum in (12.5) that merges at least two connected components of T_i, we get by the convergence in distribution of traffics (12.2) of $\mathbf{X}_i^{(N)}$

$$\frac{1}{N^{K_i-1}}\mathbb{E}\Big[\tau_N\big[\pi(T_i)(\mathbf{X}_i^{(N)})\big]\Big] = O\Big(\frac{1}{N}\Big).$$

Hence, using again Lemma 3.2, we obtain

$$\begin{aligned}\frac{1}{N^{K_i-1}}\mathbb{E}\Big[\tau_N^0\big[T_i(\mathbf{X}_i^{(N)})\big]\Big] &= \mathbb{E}\Big[\prod_{k=1}^{K_i}\sum_{\pi\in\mathcal{P}(V_{i,k})} \mu_{V_{i,k}}(\pi)\tau_N\big[\pi(T_{i,k})(\mathbf{X}_i^{(N)})\big]\Big] + O\Big(\frac{1}{N}\Big)\\ &= \mathbb{E}\Big[\tau_N^0\big[T_{i,1}(\mathbf{X}_i^{(N)})\big]\dots\tau_N^0\big[T_{i,K_i}(\mathbf{X}_i^{(N)})\big]\Big] + O\Big(\frac{1}{N}\Big)\\ &= \tau^0\big[T_{i,1}(\mathbf{x}_i)\big]\dots\tau^0\big[T_{i,K_i}(\mathbf{x}_i)\big] + o(1)\end{aligned}$$

Step 3: Contribution due to the shape of T
We have proved that

$$\begin{aligned}&\mathbb{E}\Big[\tau_N^0\big[T(\mathbf{X}_1^{(N)},\mathbf{X}_2^{(N)})\big]\Big]\\ &= \frac{(N-|V_1|)!(N-|V_2|)!}{(N-|V|)!(N-1)!}N^{K_1-1}N^{K_2-1}\Big(\prod_{i=1}^{2}\prod_{k=1}^{K_i}\tau^0\big[T_{i,k}(\mathbf{x}_i)\big]+o(1)\Big)\end{aligned}$$

Write an equivalent for the normalizing term in that expression.

$$\frac{(N-|V_1|)!(N-|V_2|)!}{(N-|V|)!(N-1)!}N^{K_1-1}N^{K_2-1}\sim N^{K_1+K_2+|V|-|V_1|-|V_2|-1}.$$

Let $\mathcal{V}$ be the set of vertices of T that belong to simultaneously to T_1 and T_2, so that $|V|-|V_1|-|V_2|=-|\mathcal{V}|$. Let $\tilde{T}=(\tilde{V},\tilde{E})$ be the undirected graph defined by

- $\tilde{V}$ is the disjoint union of $\mathcal{V}$ and of the $T_{i,k}$, $i=1,2$ and $k=1,\dots,K_i$ (recall that the later are the components of T with labels corresponding to a same family $\mathbf{X}_i^{(N)}$, $i=1,2$).
- $\tilde{E}$ is the set of ensembles $\{v,C\}$ where v is in $\mathcal{V}$ and C is a component of T such that v is a vertex of C.

It is clear that $\tilde{T}$ is a tree if and only if T is a free products of *-test graphs in the labels corresponding to $\mathbf{X}_1^{(N)}$ and of *-test graphs in the labels corresponding to $\mathbf{X}_2^{(N)}$. Assume now that T is connected. By the relation between the number of vertices and the number of edges in a graph applied to $\tilde{T}$ (Lemma 4.3), we get

$$K_1+K_2+|\mathcal{V}|\leqslant 2|\mathcal{V}|+1, \tag{12.6}$$

with equality if and only if $\tilde{T}$ is a tree. Hence, we get the expected result: for any *-test graph T,

$$\begin{aligned}\mathbb{E}\Big[\tau_N^0\big[T(\mathbf{X}_1^{(N)},\mathbf{X}_2^{(N)})\big]\Big] &= \Big(\mathbf{1}_{\big(T\text{ is a free product}\big)}+o(1)\Big)\times\Big(\prod_{i=1}^{2}\prod_{k=1}^{K_i}\tau^0\big[T_{i,k}(\mathbf{x}_i)\big]+o(1)\Big)\\ &= \tau^0\big[T(\mathbf{x}_1,\mathbf{x}_2)\big]+o(1).\end{aligned}$$

Step 4: Proof of (12.3)

Lemma 12.4. *Let $\mathbf{A}_N$ be a family of matrices and $T_1,\dots,T_n$ be *-test graphs. Let S be the *-graph obtained as the disjoint union of $T_1,\dots,T_n$. Then,*

$$\tau_N^0\big[T_1(\mathbf{A}_N)\big]\dots\tau_N^0\big[T_n(\mathbf{A}_N)\big] = \sum_\pi\frac{1}{N^{n-1}}\tau_N^0\big[\pi(S)(\mathbf{A}_N)\big],$$

where the sum is over all partitions π on V that contain at most one vertex of each T_k, $k=1,\dots,n$.

Proof of Lemma 12.4. We write $S=(V,E,\gamma,\varepsilon)$ and denote by V_k the set of vertices of T_k, $k=1,\dots,n$. Then,

$$\tau_N^0\big[T_1(\mathbf{A}_N)\big]\dots\tau_N^0\big[T_n(\mathbf{A}_N)\big]=\frac{1}{N^n}\sum_\phi\prod_{e\in E}A_{\gamma(e)}^{\varepsilon(e)}\big(\phi(e)\big),$$

where the sum is over all maps $\phi:V\to\{1,\dots,N\}$ such that $\phi_{|V_1},\dots,\phi_{|V_n}$ are injective. The sum over π in the Lemma represents all the possible situations of overlapping of the images of $\phi_{|V_1},\dots,\phi_{|V_n}$. □

Let $T_1,\dots,T_n,S$ be as in the Lemma:

$$\mathbb{E}\Big[\tau_N^0\big[T_1(\mathbf{X}_1^{(N)},\mathbf{X}_2^{(N)})\big]\dots\tau_N^0\big[T_n(\mathbf{X}_1^{(N)},\mathbf{X}_2^{(N)})\big]\Big]=\sum_\pi\frac{1}{N^{n-1}}\tau_N^0\big[\pi(S)(\mathbf{X}_1^{(N)},\mathbf{X}_2^{(N)})\big]. \tag{12.7}$$

Let π be a partition as in the sum. Denote by n_π the number of components of $\pi(S)$. If we write $T = \pi(S)$ and use the notation of the previous steps, we have to modify (12.6) into

$$K_1 + K_2 + |\mathcal{V}| \leqslant 2|\mathcal{V}| + n_\pi, \tag{12.8}$$

and obtain

$$\begin{aligned}&\mathbb{E}\Big[\tau_N^0\big[T(\mathbf{X}_1^{(N)}, \mathbf{X}_2^{(N)})\big]\Big]\\ &= \quad N^{n_\pi - 1}\Big(\mathbf{1}_{\big(\text{ the components of } T \text{ are free products }\big)} + o(1)\Big) \times \Big(\prod_{i=1}^{2}\prod_{k=1}^{K_i} \tau^0\big[T_{i,k}(\mathbf{x}_i)\big] + o(1)\Big).\end{aligned}$$

Hence, the only partition π which contributes in (12.7) is the trivial partition and we get

$$\mathbb{E}\Big[\tau_N^0\big[T_1(\mathbf{X}_1^{(N)}, \mathbf{X}_2^{(N)})\big] \dots \tau_N^0\big[T_n(\mathbf{X}_1^{(N)}, \mathbf{X}_2^{(N)})\big]\Big] \underset{N\to\infty}{\longrightarrow} \tau^0\big[T_1(\mathbf{x}_1, \mathbf{x}_2)\big] \times \dots \times \tau^0\big[T_n(\mathbf{x}_1, \mathbf{x}_2)\big].$$

□

12.2 Applications

12.2.1 Deterministic matrices

The following corollary is a direct consequence of Theorem 12.1.

Corollary 12.5. *Let $\mathbf{Y}_1^{(N)}, \dots, \mathbf{Y}_p^{(N)}$ be p families of deterministic complex matrices, and $U_1^{(N)}, \dots, U_p^{(N)}$ be independent uniform permutation matrices. Let $\mathbf{y}_1, \dots, \mathbf{y}_p$ be free families of traffics in a space of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$. Assume that for any $j = 1, \dots, p$, $\mathbf{Y}_j^{(N)}$ converges to $\mathbf{y}_j$ in distribution of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$. Then, in expectation $(U_1^{(N)}\mathbf{Y}_1^{(N)}U_1^{(N)*}, \dots, U_p^{(N)}\mathbf{Y}_p^{(N)}U_p^{(N)*})$ converges in distribution of traffics to $(\mathbf{y}_1, \dots, \mathbf{y}_p)$.*

Answer to Question 1: In particular, by Proposition 2.1, if A_N and B_N are two deterministic matrices having a limiting distribution of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$, Theorem 12.1 gives us a characterization of the limiting empirical spectral distribution of

$$H_N = A_N + U_N B_N U_N^*.$$

12.2.2 A family of operations on measures

Let U_N be an arbitrary unitary random matrix. Assume that in expectation, U_N converges to a traffic u in distribution on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$. Let μ, ν be two probability measures with compact support. Let A_N, B_N be two permutation invariant diagonal matrices such that the mean empirical eigenvalues distribution of A_N (respectively B_N) converges to μ (respectively ν). For instance, one can take for A_N a realization of the diagonal random matrix whose entries are independent, identically distributed according to μ. We deduce form Corollary 12.5 that in expectation the random matrix

$$H_N = A_N + U_N B_N U_N^*$$

has a limiting distribution of traffics $h = a + ubu^*$. In particular, the empirical eigenvalues distribution $\mathcal{L}_{H_N}$ of H_N converges in moments. Since μ and ν are compactly supported, the limiting moments characterizes of h are those of a unique probability measure. $\mathcal{L}_{H_N}$ converges to a measure that could denoted

$$\mu \boxplus_u \nu.$$

Benaych-Georges and Lévy' notion of t-freeness [5] is likely a particular case of this construction.

12.2.3 Eigenvectors of random matrices

Proposition 12.6 (A bivariate processes from the eigenvectors)**.**
Let H_N be a Hermitian random matrix. Assume that

- *H_N is invariant in law by permutation,*
- *in expectation, H_N has a limiting distribution of traffics and H_N satisfies Assumption 4) of Theorem 12.1,*
- *the limiting mean empirical eigenvalues distribution of H_N has no mass,*
- *the operator norm of H_N is uniformly bounded.*

Denote by $\big(u(i,j)\big)_{i,j=1,\dots,N}$ a matrix of eigenvectors of H_N. Then, for any r,s in $]0,1[$, the quantity

$$B_N(r,s) = \frac{1}{N}\sum_{i=1}^{\lfloor rN\rfloor}\sum_{j=1}^{\lfloor sN\rfloor}|u(i,j)|^2 \tag{12.9}$$

converges in expectation as N goes to infinity.

Proof. Consider the diagonal matrix

$$J_r = \mathrm{diag}\ (\underbrace{1,\dots,1}_{\lfloor rN\rfloor},0,\dots,0).$$

By Theorem 12.1, in expectation (H_N, J_r) has a limiting distribution of traffics. In particular, for any polynomial P, the quantity

$$\frac{1}{N}\mathrm{Tr}\big[P(H_N)\circ J_r\big]$$

converges in expectation as N goes to infinity. Let q_r be the quantile of order r of the limiting eigenvalues distribution of H_N:

$$q_r = \inf\Big\{t\in\mathbb{R}\Big|\mu(]-\infty,r])\geqslant x\Big\}.$$

Denote by f the indicator function of the interval $]-\infty,q_r]$. Let $\varepsilon>0$ and P be a polynomial ε-closed to f in uniform norm to f on $[-B,B]$, where $\|H_N\|\leqslant B$. Then,

$$\begin{aligned}
&\Bigg|\mathbb{E}\Bigg[\frac{1}{N}\sum_{i=1}^{\lfloor rN\rfloor}\sum_{j=1}^{\lfloor sN\rfloor}|u(i,j)|^2-\frac{1}{N}\mathrm{Tr}\big[P(H_N)\circ J_r\big]\Bigg]\Bigg|\\
&\leqslant\ \Bigg|\mathbb{E}\Bigg[\frac{1}{N}\sum_{i=1}^{N}f(\lambda_i)\sum_{j=1}^{\lfloor sN\rfloor}|u(i,j)|^2-\frac{1}{N}\mathrm{Tr}\big[P(H_N)\circ J_s\big]\Bigg]\Bigg|\\
&\quad+\Bigg|\mathbb{E}\Bigg[\frac{1}{N}\sum_{i=1}^{\lfloor rN\rfloor}\sum_{j=1}^{\lfloor sN\rfloor}|u(i,j)|^2-\frac{1}{N}\sum_{i=1}^{N}f(\lambda_i)\sum_{j=1}^{\lfloor sN\rfloor}|u(i,j)|^2\Bigg]\Bigg|.
\end{aligned}$$

On the one hand, one has

$$\begin{aligned}
&\Bigg|\mathbb{E}\Bigg[\frac{1}{N}\sum_{i=1}^{N}f(\lambda_i)\sum_{j=1}^{\lfloor sN\rfloor}|u(i,j)|^2-\frac{1}{N}\mathrm{Tr}\big[P(H_N)\circ J_s\big]\Bigg]\Bigg|\\
&\leqslant\ \frac{1}{N}\sum_{i=1}^{N}\Big|\mathbb{E}\big[f(\lambda_i)-P(\lambda_i)\big]\Big|\,\Big|\sum_{j=1}^{\lfloor sN\rfloor}|u(i,j)|^2\leqslant\varepsilon.
\end{aligned}$$

Moreover, by [19, Lemma 9.1], there exists η arbitrary small such that

$$\mathbb{E}\big[\lambda_{1+\lfloor(r-\eta)N\rfloor}\big]\underset{N\to\infty}{\longrightarrow}q_{r-\eta}$$
$$\mathbb{E}\big[\lambda_{1+\lfloor(r+\eta)N\rfloor}\big]\underset{N\to\infty}{\longrightarrow}q_{r+\eta}.$$

Since the mean limiting eigenvalues distribution of H_N has no mass, one has $q_{r-\eta} < q_r < q_{r+\eta}$. Hence, for N large enough, we get

$$\begin{aligned}
&\left|\mathbb{E}\left[\frac{1}{N}\sum_{i=1}^{\lfloor rN\rfloor}\sum_{j=1}^{\lfloor sN\rfloor}|u(i,j)|^2 - \frac{1}{N}\sum_{i=1}^{N} f(\lambda_i)\sum_{j=1}^{\lfloor sN\rfloor}|u(i,j)|^2\right]\right| \\
= \quad & \left|\mathbb{E}\left[\frac{1}{N}\sum_{i=\lfloor(r-\eta)N\rfloor}^{\lfloor rN\rfloor}\sum_{j=1}^{\lfloor sN\rfloor}|u(i,j)|^2 - \frac{1}{N}\sum_{i=\lfloor(r-\eta)N\rfloor}^{\lfloor(r+\eta)N\rfloor} f(\lambda_i)\sum_{j=1}^{\lfloor sN\rfloor}|u(i,j)|^2\right]\right| \\
\leqslant \quad & 4\eta.
\end{aligned}$$

As η and ε go to zero, we obtain the expected result.

□

12.2.4 Wigner matrices

Corollary 12.7. *Let $\mathbf{X}_N = (X_1, \dots, X_p)$ be a family of independent real or complex Wigner matrices. Let $\mathbf{Y}_N$ be a family of complex deterministic matrices having a limiting distribution of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle$. Then, in expectation $(\mathbf{X}_N, \mathbf{Y}_N)$ converge on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle$ to a family of traffics $(x_1, \dots, x_p, \mathbf{y})$ such that $x_1, \dots, x_p, \mathbf{y}$ are free.*

Let X_N be a real or complex Wigner matrix as in Proposition 4.2. Then, X_N satisfies the assumption *2)* of Theorem 12.1, the independence by permutation, since its diagonal entries and its sub-diagonal entries respectively are independent and identically distributed. The assumption *4)*, is a slight generalization of Proposition 4.2.

Lemma 12.8. *The matrix X_N satisfies (12.2).*

Proof. Let $T_1, \dots, T_n$ be test graphs in one variable, and denote by T the graph obtained as the disjoint union of $T_1, \dots, T_n$. By Lemma 12.4,

$$\mathbb{E}\Big[\tau_N^0\big[T_1(X_N)\big]\dots\tau_N^0\big[T_n(X_N)\big]\Big] = \ = \sum_{\pi}\frac{1}{N^{n-1}}\mathbb{E}\Big[\tau_N^0\big[\pi(T)(X_N)\big]\Big],$$

where the sum is as in the Lemma. For any such a partition π, denote by $T_1^\pi, \dots, \tilde{T}_{m_\pi}^\pi$ the connected components of $\pi(T)$. By the independence of the entries of X_N,

$$\begin{aligned}
&\mathbb{E}\Big[\tau_N^0\big[T_1(X_N)\big]\dots\tau_N^0\big[T_n(X_N)\big]\Big] \\
&\quad = \sum_{\pi}\frac{N^{m_\pi}}{N^n}\mathbb{E}\Big[\tau_N^0\big[T_1^\pi(X_N)\big]\Big]\dots\mathbb{E}\Big[\tau_N^0\big[T_{m_\pi}^\pi(X_N)\big]\Big],
\end{aligned}$$

Each expectation converges as N goes to infinity. We always has $m_\pi \leqslant n$, expect for the trivial partition. Hence, we get

$$\mathbb{E}\Big[\tau_N^0\big[T_1(X_N)\big]\dots\tau_N^0\big[T_n(X_N)\big]\Big] \underset{N\to\infty}{\longrightarrow} \tau_x^0[T_1]\dots\tau_x^0[T_n],$$

where τ_x is the mean limiting distribution of traffics of X_N. □

Proof of Corollary 12.7. Let U_N be a uniform permutation matrix, independent of $\mathbf{X}_N$. By the invariance by permutation of a Wigner matrix, the families $(\mathbf{X}_N, \mathbf{Y}_N)$ and $(\mathbf{X}_N, U_N\mathbf{Y}_N U_N^*)$ have the same mean distribution of traffics. Hence, Theorem 12.1 can be applied to the families $(X_1), \dots, (X_p), \mathbf{Y}_N$. □

12.2.5 Permutation Matrices

Corollary 12.9. *Let $\mathbf{U}_N = (U_1, \dots, U_p)$ be a family of independent permutation matrices. Let $\mathbf{Y}_N$ be a family of complex deterministic matrices having a limiting distribution of traffics on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle$. Then, in expectation $(\mathbf{U}_N, \mathbf{Y}_N)$ converge on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle$ to a family of traffics $(u_1, \dots, u_p, \mathbf{y})$ such that $u_1, \dots, u_p, \mathbf{y}$ are free.*

In particular, a couple of independent permutation matrices has in expectation a limiting distribution of traffics τ, such that τ^0 is the indicator function of free products of directed lines, as in Figure 15.

Let U_N be a real or complex uniform permutation matrix. Then, U_N satisfies the assumption *2)* of Theorem 12.1, the independence by permutation. It remains to prove assumption *4)*, which is a slight generalization of Proposition 4.5.

Lemma 12.10. *The matrix U_N satisfies (12.2).*

Proof. Real case: Let $T = (V, E)$ be test graphs in one variable whose directed edges are of multiplicity one. We have seen in the proof of Proposition 4.5 that it is sufficient to consider such test graphs. We have shown that $\tau_N^0\big[T(U_N)\big]$ is possibly nonzero only if T is a test graph T_K^c (closed path) or T_K^o (open path) for a certain positive integer K. Let σ_N be the permutation of $\{1, \dots, N\}$ associated to U_N. Then, for any $K_1, \dots, K_n, L_1, \dots, L_m$, the number

$$\mathbb{E}\Big[\tau_N^0\big[T_{K_1}^o(U_N)\big] \dots \tau_N^0\big[T_{K_n}^o(U_N)\big] \tau_N^0\big[T_{L_1}^c(U_N)\big] \dots \tau_N^0\big[T_{L_m}^c(U_N)\big]\Big]$$

is the probability that, choosing $i_1, \dots, i_n, j_1, \dots, j_m$ uniformly and independently on $\{1, \dots, N\}$ one has

- i_k belongs to a cycle of length K_k of σ_N for any $k = 1, \dots, n$,
- j_k belongs to a cycle of length bigger than L_k of σ_N for any $k = 1, \dots, m$.

This probability tends to zero or one, depending if n is positive or not, which correspond to the limit of

$$\mathbb{E}\Big[\tau_N^0\big[T_{K_1}^o(U_N)\big]\Big] \dots \mathbb{E}\Big[\tau_N^0\big[T_{K_n}^o(U_N)\big]\Big] \mathbb{E}\Big[\tau_N^0\big[T_{L_1}^c(U_N)\big]\Big] \dots \mathbb{E}\Big[\tau_N^0\big[T_{L_m}^c(U_N)\big]\Big].$$

Complex case: The proof is still based on the Hadamard structure of a complex permutation matrix. Let $T_1, \dots, T_n$ be *-test graphs in one variable. Let $T = (V, E, \varepsilon)$ be the *-graph in one variable obtained as the disjoint union of $T_1, \dots, T_n$. By Lemma 12.4,

$$\mathbb{E}\Big[\tau_N^0\big[T_1(U_N)\big] \dots \tau_N^0\big[T_n(U_N)\big]\Big] = \sum_{\pi} \frac{1}{N^{n-1}} \mathbb{E}\Big[\tau_N^0\big[\pi(T)(U_N)\big]\Big],$$

where the sum is over all partition π on V that contains at most one element of each V_k, $k = 1, \dots, n$. For any such a partition π, denote by $T_1^\pi, \dots, \tilde{T}_{m_\pi}^\pi$ the connected components of $\pi(T)$. We write, as in the proof of Proposition 4.5, $U_N = V_N \circ M_N$. Then, we get

$$\begin{aligned}
&\mathbb{E}\Big[\tau_N^0\big[T_1(U_N)\big] \dots \tau_N^0\big[T_n(U_N)\big]\Big] \\
&= \sum_{\pi} \frac{1}{N^n} \sum_{\substack{\phi:\pi(V) \to \{1,\dots,N\} \\ \text{injective}}} \mathbb{E}\Big[\prod_{e \in \pi(E)} V_N^{\pi(\varepsilon)(e)}\big(\phi(e)\big)\Big] \mathbb{E}\Big[\prod_{e \in \pi(E)} M_N^{\pi(\varepsilon)(e)}\big(\phi(e)\big)\Big] \\
&= \sum_{\pi} \mathbf{1}_{\pi(T) \text{ is Hermitian}} \frac{1}{N^n} \sum_{\substack{\phi:\pi(V) \to \{1,\dots,N\} \\ \text{injective}}} \mathbb{E}\Big[\prod_{e \in \pi(E)} V_N^{\pi(\varepsilon)(e)}\big(\phi(e)\big)\Big]
\end{aligned}$$

For any partition π as in the sum, denote by $T_1^\pi, \dots, \tilde{T}_{m_\pi}^\pi$ the connected components of $\pi(T)$. Then,

$$\begin{aligned}
&\mathbb{E}\Big[\tau_N^0\big[T_1(U_N)\big] \dots \tau_N^0\big[T_n(U_N)\big]\Big] \\
&\quad = \sum_{\pi} \mathbf{1}_{\pi(T) \text{ is Hermitian}} \frac{N^{m_\pi}}{N^n} \mathbb{E}\Big[\tau_N^0\big[T_1^\pi(V_N)\big] \dots \tau_N^0\big[T_{m_\pi}^\pi(V_N)\big]\Big].
\end{aligned}$$

By the real case, for any π,

$$\mathbb{E}\Big[\tau_N^0\big[T_1^\pi(V_N)\big] \dots \tau_N^0\big[T_{m_\pi}^\pi(V_N)\big]\Big] \tau_v^0[T_1^\pi] \dots \tau_v^0[T_{m_\pi}^\pi] + o(1),$$

where τ_v is the mean limiting distribution of traffics of V_N. Hence, the only terms that contributes is when π is the trivial partition. In that case, $\pi(T)$ Hermitian means that each T_k is Hermitian, $k = 1, \dots, n$, so

$$\begin{aligned}
&\mathbb{E}\Big[\tau_N^0\big[T_1(U_N)\big]\dots\tau_N^0\big[T_n(U_N)\big]\Big]\\
&= \mathbf{1}_{T_1 \text{ is Hermitian}}\tau_v^0[T_1^\pi]\dots\mathbf{1}_{T_n \text{ is Hermitian}}\tau_v^0[T_{m_\pi}^\pi] + o(1)\\
&= \tau_u^0[T_1^\pi]\dots\tau_u^0[T_{m_\pi}^\pi] + o(1).
\end{aligned}$$

where τ_u is the mean limiting distribution of traffics of U_N.

□

13 A central limit theorem for the sum of free traffics

Let $\mathbf{x} = (x_n)_{n\geqslant 1}$ be a sequence of identically distributed self-adjoint free traffics in a space of traffics $(\mathcal{A}, .^*, \tau)$. We are interested in the limiting behavior of

$$m_n = \frac{x_1 + \dots + x_n}{\sqrt{n}}, \text{ as } n \text{ goes to infinity.}$$

If $x_1, \dots, x_n$ were independent, centered, real random variables with finite variance, Laplace's central limit theorem states that m_n converges in law to a Gaussian random variable. If $x_1, \dots, x_n$ were free, centered, self-adjoint non commutative random variables, Voiculescu's central limit theorem [23] states that m_n converges in *-distribution to a semicircular non commutative random variable. The case of free traffics turns out to be an interpolation of these two situations.

Assume that the traffics $x_1, \dots, x_n$ are centered, i.e. $\Phi^{(\tau)}(x_1) = 1$, and of variance one, i.e. $\Phi^{(\tau)}(xx^*) = \Phi^{(\tau)}(x^2) = 1$. By fixing these parameters, the limit in law or in *-distribution of m_n is completely determined in the classical and the free context. This is not the case for traffics, where we can read in the common distribution of $x_1, \dots, x_n$ a parameter that heuristically measure how the traffics are closed to the classical or the free cases.

We split the variance of x_1 into two parts

$$\tau^0\big[T_1(x_1)\big] = p, \quad \tau^0\big[T_2(x_1)\big] = (1-p),$$

where

- T_1 is the test graph with one vertex and two edges labelled x,
- T_2 is the test graph with two vertices i and j and two edges labelled x, one from i to j and the other one from j to i.

(We have rightly $\Phi^{(\tau)}(x^2) = \tau\big[T_2(x_1)\big] = \tau^0\big[T_1(x_1)\big] + \tau^0\big[T_2(x_1)\big]$).

Equivalently,

$$p = \Phi^{(\tau)}(\Delta_x^2), \quad (1-p) = \Phi^{(\tau)}\big((x-\Delta_x)^2\big)$$

(we have rightly $\Phi^{(\tau)}(x\Delta_x) = \Phi^{(\tau)}(\Delta_x x) = \Phi^{(\tau)}(\Delta_x^2)$).

Theorem 13.1 (Central limit theorem).
With the notations above, the sequence of traffics $(m_n)_{n\geqslant 1}$ converges in distribution on $\mathcal{G}_{cyc}\langle \mathbf{x}, \mathbf{x}^*\rangle$ to the traffic

$$m = \sqrt{p}\; d + \sqrt{1-p}\; s,$$

where

1. *the law of d is the diagonal distribution associated to the standard Gaussian measure μ on $\mathbb{R}$ (Definition 4.6):*

$$d\mu(x) = \frac{1}{\sqrt{2\pi}}\exp\Big(-\frac{x^2}{2}\Big)dx,$$

2. *the law of s is the limiting distribution of traffics of a real or complex Wigner matrix as in Proposition 4.2.*
3. *the traffics d and s are free.*

From Theorem 13.1, one can recover the classical Lagrange's central limit theorem for random variable having all their moments. This fact is straightforward since we can lift commutative random variables into diagonal traffics, and then reformulate the classical central limit theorem in terms of these traffics.

From Theorem 13.1, we can also recover Voiculescu's central limit theorem. Indeed, by [24, Theorem 6.1] and minor modification of Proposition 4.2, a random unitary matrix U_N distributed according to the Haar measure on the unitary group has a limit u in mean distribution of traffics. Let x be a self adjoint non commutative random variable in a *-probability space $(\mathcal{A}, .^*, \Phi)$. We can lift x into a traffic t_x by considering a diagonal traffic d_x whose spectrum is the one of x, free from u, and setting $t_x = u d_x u^*$. The non commutative random variable induced by t_x is well x, and a free family of traffics $t_1, \dots, t_n$ distributed as t_x induces a *-free family of variables $x_1, \dots, x_n$, distributed as x (by Voiculescu's asymptotic freeness theorem in [2]). We do not known how to lift an arbitrary family of non commutative random variables.

Remarks:

- The support of the injective version of the distribution of $\sqrt{p}\, d + \sqrt{1-p}\, s$ consists of flowery double trees, as in Figure 15 but in only one color.
- A matrix model for the traffic $m = \sqrt{p}\, d + \sqrt{1-p}\, s$ is the sum $M_N = \sqrt{p}\, D_N + \sqrt{1-p}\, S_N$, where S_N is a standard Wigner matrix, independent of diagonal matrix D_N whose entries are independent standard Gaussian variables. It is known by Voiculescu's asymptotic freeness theorem [2], that the limiting spectral distribution of M_N is the free convolution of the centered Gaussian and the semicircular distribution, weights $\sqrt{p}$ and $\sqrt{1-p}$ determining the variance. In that case, the *-non commutative random variable induced by d and s are *-free (recall that in general, the traffic-freeness of the limiting *-distribution of matrices does not imply their *-freeness).
- From the author's knowledge, the only appearances of free convolution (in the sense of Voiculescu) of gaussian and semicircular variables in the literature concern the limiting eigenvalues distribution of Laplacian matrices of random graphs [9, 12]. See also [7, 13] for related questions.
- The convergence of m_n actually holds on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ (see the proof), but we do not know how to characterize its limiting distribution in elegant terms.

Proof. Since the traffics are self-adjoint, it is sufficient to consider test graph (and not *-test graphs). Let $T = (V, E)$ be a cyclic test graph in one variable. By the rule of multi-linearity for τ, Formulae (7.5) and (7.6),

$$\tau^0\big[T(m_n)\big] = \frac{1}{n^{\frac{|E|}{2}}} \sum_{\tilde T = (V,E,\gamma)} \tau^0\big[\tilde T(\mathbf{x})\big],$$

where the sum is over all maps $\gamma : E \to \{1, \dots, n\}$. Let π be a partition of E. We denote by $\Gamma_\pi^{(n)}$ the set of maps $\gamma : E \to \{1, \dots, n\}$ such that $\gamma(e) = \gamma(e')$ if and only if e and e' belong to the same block of π. Since the traffics $x_1, \dots, x_n$ are identically distributed, for any $\tilde T = (V, E, \gamma)$ as in the sum, the number $\tau^0\big[\tilde T(\mathbf{x})\big]$ depends only on the partition π such that $\gamma \in \Gamma_\pi^{(n)}$. We denote this number by a_π. Hence, we get

$$\tau^0\big[T(m_n)\big] = \frac{1}{n^{\frac{|E|}{2}}} \sum_{\pi \in \mathcal{P}(E)} a_\pi \times \mathrm{Card}\ \big(\Gamma_\pi^{(n)}\big).$$

For any π in $\mathcal{P}(E)$, denote by $|\pi|$ its number of blocks. Then, Card $\big(\Gamma_\pi^{(n)}\big) = n \times (n-1) \times \cdots \times (n - |\pi| + 1) \sim n^{|\pi|}$.

If π possesses a block of cardinal one, we claim that $a_\pi = 0$. Indeed, let $\gamma \in \Gamma_\pi^{(n)}$ and denote $\tilde T = (V, E, \gamma, \varepsilon)$. Let n_0 in $\{1, \dots, n\}$ appearing once as a label of T. By the freeness of the traffics $x_1, \dots, x_n$, one has $\tau^0\big[\tilde T(\mathbf{x})\big]$ if this edge is not a loop (otherwise, since $\tilde T$ is cyclic, it is never a free product of test graphs). Nevertheless, if this edge is a loop, then we can factorizes $\tau^0[x_{n_0}]$ in the computation of $\tau^0\big[\tilde T(\mathbf{x})\big]$, which is zero. This proves the claim. We then get that if $|\pi| > \frac{|E|}{2}$ or $|\pi| = \frac{|E|}{2}$ and π is not a pair partition (each block of π is of cardinal two), then $a_\pi = 0$.

Hence, if we denote by $\mathcal{P}_2(E)$ the set of pair partitions of E, we get

$$\tau^0\big[T(m_n)\big] \quad = \quad \sum_{\pi \in \mathcal{P}_2(E)} a_\pi + o(1).$$

Let π be in $\mathcal{P}_2(E)$ and assume $a_\pi \neq 0$. Let e be an edge of $\tilde T$. By the same reasoning as above, the other edge e' with the same label must share the same vertices as e, and if e is not a loop, the cyclic nature of $\tilde T$ forces e and e' to have the opposite directios.

Hence $\tilde T$ is a free product of cyclic test graphs that are either double loops (one vertex and two edges) or double arrows (two vertices and two edges joining this vertices in opposite directions). All these elementary test graphs are labelled by different labels. To sum up, the graph of $\tilde T$ consists in a double tree T_0 with flowers $F_1, \dots, F_K$ attached at its vertices. Each flower must have an even number of petals, and the partition π must gather twin edges of T_0 and pair of petals attached at a same flower. Denote by $2m_k$ the number of petals of F_k, $k = 1, \dots, K$. By Lemma 4.3 that gives the relation between the number of vertices and edges in a tree, the number of edges of T_0 is $2(K-1)$. We get

$$\tau^0\big[T(m_n)\big] \quad = \quad (1-p)^{K-1} \prod_{k=1}^{K} p^{m_k} \text{Card } \mathcal{P}_2(2m_k),$$

where $\mathcal{P}_2(2m)$ denotes the set of pair partitions of $2m$ elements. But

$$\text{Card } \mathcal{P}_2(2m) = (2m-1) \times (2m-3) \dots 5 \times 3 \times 1 = \mathbb{E}[X^{2m}]$$

where X is a random variable distributed according to the standard gaussian measure (by a basic enumeration and by integration by part respectively).

Now, let d and s be as in the Theorem. For any cyclic test graph $T = (V, E)$ in one variable, by the rule of the sum for τ, Formula (7.6),

$$\tau^0\big[T(\sqrt{p}d + \sqrt{1-p}s)\big] \quad = \quad \sum_{\tilde T = (V,E,\gamma)} \tau^0\big[\tilde T(\sqrt{p}d, \sqrt{1-p}s)\big],$$

where the sum is over all maps $\gamma : E \to \{1, 2\}$. By the definition of freeness of traffics, the support of the injective version of the distribution of (d, s) consists of free products of double trees and flowers. If T is such a test graph and is as above with the notations T_0, $F_1, \dots, F_K$, the only map γ which makes $\tau^0\big[\tilde T(\sqrt{p}d, \sqrt{1-p}s)\big]$ possibly non zero consists of labeling the edge of T_0 with labels corresponding to s and the edges of the flowers by the one corresponding to d. By the rule for the homogeneity for τ, Formula (7.5), we get

$$\tau^0\big[T(\sqrt{p}d + \sqrt{1-p}s)\big] \quad = \quad (1-p)^{K-1} \prod_{k=1}^{K} p^{m_k} \mathbb{E}[X^{2m_k}]$$

as expected. □

Part V

The distribution of traffics of graphs with uniformly bounded degree

Let $\mathbf{G}_N = (G_1^{(N)}, \dots, G_p^{(N)})$ be a family of directed graphs with no multiple edges nor loops, sharing the same N vertices $v_1, \dots, v_N$. We arbitrary label these vertices by the integers $\{1, \dots, N\}$. The following family of matrices $\mathbf{A}_N = (A_1^{(N)}, \dots, A_p^{(N)})$, defined up to this labeling, is called an adjacency family of matrices of $\mathbf{G}_N$. For any $j = 1, \dots, p$, $A_j^{(N)} = \big(A_j^{(N)}(m,n) \big)_{m,n=1,\dots,N}$ is the N by N (0,1)-matrix given by: for any $m, n = 1, \dots, N$,

$$A_j^{(N)}(m,n) = \left\{ \begin{array}{ll} 1 & \text{if } (m,n) \text{ is an edge of } G_N, \\ 0 & \qquad\quad \text{otherwise.} \end{array} \right.$$

The distribution of traffics $\tau_{\mathbf{G}_N}$ of $\mathbf{G}_N$ is defined as the distribution of $\mathbf{A}_N$ and does not depend on the labeling of vertices of $\mathbf{G}_N$. Remark that $\mathbf{G}_N$ can be seen as a colored graph, or equivalently as a graph whose edges are labeled by indeterminates $x_1, \dots, x_p$. In other words, $\mathbf{G}_N$ is a test graph in p variables $\mathbf{x}$. It should be denoted that since A_N is a (0,1)-matrix, we know $\tau_{\mathbf{G}_N}$ on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^* \rangle$ as soon as we know it on $\mathbf{G}_{cyc}\langle \mathbf{x} \rangle$.

Infinite rooted random graphs with uniformly bounded degree constitute a model for the limit of large graphs for the weak local convergence [6, 1]. We define the formal distribution of traffics of such infinite graphs (we say simply distribution of traffics, even if we have not verified that such distributions are positive). We show that the distribution of traffics of such graphs characterizes the law of the graph. We prove that the convergence in distribution of traffics of $\mathbf{G}_N$ is equivalent to its weak local convergence. We interpret the free product in that context.

Preliminary definitions: random rooted graphs

Locally finite, colored, rooted graphs

Denote by $\mathcal{G}^\infty\langle \mathbf{x} \rangle$ the set, up to isomorphism, of all directed graphs, without multiple edges nor loops, whose degree of each vertex is finite and whose edges are labelled by an element of the family of labels (or colors) $\mathbf{x}$. We denote by $\mathcal{G}^\infty_*\langle \mathbf{x} \rangle$ the set, up to isomorphism, of rooted elements of $\mathcal{G}^\infty\langle \mathbf{x} \rangle$, i.e. couples (G, v) where G is in $\mathcal{G}^\infty\langle \mathbf{x} \rangle$ and v is a vertex of G, called its root.

When we say up to isomorphism, we mean the following. Let $g_1 = (G_1, v_1)$ and $g_2 = (G_2, v_2)$ be such rooted graphs. Then, (the isomorphism class of) g_1 and g_2 are equal (respectively $g_1 \leqslant g_2$) whenever there exists an isomorphism (respectively an injection) of oriented colored graphs $\phi : G_1 \to G_2$ such that $\phi(v_1) = \phi(v_2)$.

A topology on $\mathcal{G}^\infty_*\langle \mathbf{x} \rangle$

For any integer $p \geqslant 0$, define $\mathcal{G}^p_*\langle \mathbf{x} \rangle$ as the set of all graphs in $\mathcal{G}^\infty_*\langle \mathbf{x} \rangle$ whose vertices are at distance at most p of the root. For any integer $p \geqslant 0$ and any (G, v) in $\mathcal{G}^\infty_*\langle \mathbf{x} \rangle$, we denote by $(G, v)_{|p}$ in $\mathcal{G}^p_*\langle \mathbf{x} \rangle$ the truncation of (G, v) of depth p, i.e. the connected sub-graph of G, rooted at v, constituted by the vertices of G that are at distance at most p of v and by the edges linking these vertices.

For any integer $p \geqslant 1$ and any rooted graph (H, w) in $\mathcal{G}^p_*\langle \mathbf{x} \rangle$, we set

$$O_{p,(H,w)} = \Big\{ \ (G, v) \in \mathcal{G}^\infty_*\langle \mathbf{x} \rangle \ \Big| \ (G, v)_{|p} = (H, w) \ \Big\}.$$

The set $\mathcal{G}^\infty_*\langle \mathbf{x} \rangle$ is equipped with the topology generated by the collections of all $O_{p,(H,w)}$, $p \geqslant 1$ and (H, w) in $\mathcal{G}^p_*\langle \mathbf{x} \rangle$ (this topology is actually metrizable by an ultrametric distance [1]). In particular, $\mathcal{G}^\infty_*\langle \mathbf{x} \rangle$ is the closure of the union of the $\mathcal{G}^p_*\langle \mathbf{x} \rangle$, $p \geqslant 1$.

Random graphs

A random element of $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$ is map $\Omega \to \mathcal{G}_*$, where Ω is a probability space, measurable for the Borel algebra on $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$. We often simply say that (G,v) in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$ is a random graph, instead of saying that $\omega \mapsto \big(G(\omega), v(\omega)\big)$ is a measurable map $\Omega \to \mathcal{G}_*$. Such a random graph is referred as a locally finite, rooted, random graph.

Definition 13.2 (Law of random graphs).
The law of (G,v) in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$ is the knowledge of $\mathbb{P}\big((G,v)_p = (H,w)\big)$ for any integer $p \geqslant 0$ and any graph (H,w) in $\mathcal{G}_*^p\langle \mathbf{x}\rangle$.

14 Distribution of traffics of locally finite, rooted, random graphs

Definition 14.1 (Rooted trace of test graphs).
Let (G,v) in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$ be a random graph. Let (T,r) be a test graph in $\mathcal{G}\langle \mathbf{x}\rangle$ with a distinguished vertex r, called its root. We denote by $\tau\big[(T,r)(G,v)\big]$ the number of homomorphisms of T into G which send r on v (we take into account the directions and the colors of the edges of T).

Definition 14.2 (Integrability of random graphs).
A random graph (G,v) in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$ is said to be integrable whenever, for any test graph T in $\mathcal{G}\langle \mathbf{x}\rangle$ and any vertex r of T, the expectation of $\tau\big[(T,r)(G,v)\big]$ is finite.

These numbers can be infinite since we can find (T,r) such that $\tau\big[(T,r)(G,v)\big]$ is the expected degree of the root.

Definition 14.3 (Traffic-stationary random graphs).
Let (G,v) in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$ be an integrable random graph. It is said to be traffic-stationary whenever for any (T,r), the expectation of $\tau\big[(T,r)(G,v)\big]$ does not depend on r.

Definition 14.4 (Distribution of traffics of random rooted graphs).
Let $g = (G,v)$ in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$ be an integrable, traffic-stationary, random graph. The mean distribution of traffic of g is the map

$$\begin{array}{cccc} \tau_g: & \mathcal{G}\langle \mathbf{x}\rangle & \to & \mathbb{C} \\ & T & \mapsto & \tau_g[T] := \mathbb{E}\Big[\tau\big[T(g)\big]\Big], \end{array}$$

where $\tau\big[T(g)\big]$ is the common value of $\tau\big[(T,r)(G,v)\big]$ for any choice of vertex r of T.

The injective version τ_g^0 of the distribution of traffics τ_g of g is given by the expectation of the common value of $\tau^0\big[(T,r)(G,v)\big]$ for any choice of vertex r of T, where $\tau^0\big[(T,r)(G,v)\big]$ is the number of injective maps form the set of vertices of T to the set of vertices of G_N, sending r to v, that are homomorphisms of colored graphs.

Proposition 14.5 (Consistency of definitions).
Let G_N in $\mathcal{G}^\infty\langle \mathbf{x}\rangle$ be a finite random graph with N vertices. Conditionally on G_N, let v_N be a vertex of G_N chosen uniformly. Then, (G_N, v_N) is integrable and traffic-stationary. Moreover, the distribution of traffic of an adjacency matrix of G_N is the distribution of traffic of (G_N, v_N) in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$.

15 Random graph with uniformly bounded degree.

Let (G,v) in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$ be a random graph. It is said to be uniformly bounded whenever there exists $K \geqslant 1$ such that almost surely the degrees of the vertices of G are less than K. We denote by $\mathcal{G}_*^{bnd}\langle \mathbf{x}\rangle$ the set of random graphs in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$, integrable, traffic-stationary and uniformly bounded.

A sequence of random graph in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$ is said to be uniformly bounded whenever all the graphs are uniformly bounded with the same constant K.

15.1 Characterization of the law

Proposition 15.1 (The distribution of traffics characterizes the law of the graph).
Two random graphs in $\mathcal{G}_*^{bnd}\langle \mathbf{x}\rangle$ have the same distribution of traffics if and only if they have the same law of random graphs.

Proof. Let $g=(G,v)$ in $\mathcal{G}_*^{bnd}\langle \mathbf{x}\rangle$. Its distribution of traffic τ_g is defined by its law (we do not need boundedness, see (15.1) below) and to show the reciprocal, it is sufficient to show that we can compute $\mathbb{P}\big((G,v)_p=(T,r)\big)$ in terms of τ_g, for any integer $p\geqslant 0$ and any graph (T,r) in $\mathcal{G}_*^{p}\langle \mathbf{x}\rangle$. This is done by an exclusion-inclusion principle.

For any (T,r) in $\mathcal{G}_*^{p}\langle \mathbf{x}\rangle$, and any integer $p\geqslant 1$, one has

$$\begin{aligned}&\mathbb{E}\Big[\tau^0\big[(T,r)(G,v)\big]\Big]\\ &\quad=\sum_{\substack{(H,w)\in\mathcal{G}_*^p\langle\mathbf{x}\rangle\\(H,w)\geqslant(T,r)}}\tau^0\big[(T,r)(H,w)\big]\times\mathbb{P}\big((G,v)_p=(H,w)\big).\end{aligned}\tag{15.1}$$

The symbol $(H,w)\geqslant(T,r)$ means that T is a subgraph of H up to an isomorphism of colored, oriented, rooted graph. But the set $\mathcal{G}_*^{p}\langle \mathbf{x}\rangle$ equipped with the order relation $\geqslant$ is a finite partially ordered set (in short, a poset). Hence, we get

$$\begin{aligned}&\mathbb{P}\big((G,v)_p=(T,r)\big)\\ &\quad=\frac{1}{\tau^0\big[(T,r)(T,r)\big]}\sum_{\substack{(H,w)\in\mathcal{G}_*^p\langle\mathbf{x}\rangle\\(H,w)\geqslant(T,r)}}\mathbb{E}\Big[\tau^0\big[(H,w)(G,v)\big]\Big]\times\mu_p\big((H,w),(T,r)\big),\end{aligned}\tag{15.2}$$

where μ_p is the Möbius map of the poset $\mathcal{G}_*^{p}\langle \mathbf{x}\rangle$ (see [23]).

□

15.2 Weak local convergence

Definition 15.2 (The weak local convergence of finite graphs [6, 1]).
Let $\big((G_N,v_N)\big)_{N\geqslant 1}$ be a sequence of random graphs in $\mathcal{G}_*^{\infty}\langle \mathbf{x}\rangle$ and (G,v) in $\mathcal{G}_*^{\infty}\langle \mathbf{x}\rangle$ random. We say that (G_N,v_N) converges weakly locally to (G,v) whenever for any integer $p\geqslant 1$ and any (H,w) in $\mathcal{G}_*^{p}\langle \mathbf{x}\rangle$, one has

$$\mathbb{P}\big((G_N,v_N)_p=(H,w)\big)\underset{N\to\infty}{\longrightarrow}\mathbb{P}\big((G,v)_p=(H,w)\big).\tag{15.3}$$

If G_N is a sequence of finite graphs, we say that it convergences weakly locally to (G,v) whenever (G_N,v_N) converges weakly locally to (G,v), where v_N is chosen uniformly on the vertices of G_N.

Theorem 15.3 (Weak local convergence and convergence in distribution of traffics).
Let $(G_N)_{N\geqslant 1}$ be a uniformly bounded sequence of random graphs. Then, in expectation G_N has a limiting distribution of traffics τ if and only if it weakly locally converges to a random graph $g=(G,v)$ in $\mathcal{G}_*^{\infty}\langle \mathbf{x}\rangle$. In this case, g is integrable and traffic-stationary, and τ is the distribution of traffics of g.

Proof. Let G_N be a finite graph in $\mathcal{G}$ and v_N be a random vertex of G_N chosen uniformly. Then (G_N,v_N) is in $\mathcal{G}_*^{bnd}\langle \mathbf{x}\rangle$. In particular, Formulae (15.1) and (15.2) hold. Hence, if G_N converges weakly locally to g, then g is also in $\mathcal{G}_*^{bnd}\langle \mathbf{x}\rangle$. We get by (15.1) that for any test graph T in one variable and any vertex r of T, one has

$$\tau^0\big[T(g)\big]:=\lim_N\mathbb{E}\Big[\tau^0\big[(T,r)(G_N,v_N)\big]\Big]\text{ exists,}$$

and does not depend on r. Since $\mathbb{E}\Big[\tau^0\big[(T,r)(G_N,v_N)\big]\Big]=\mathbb{E}\Big[\tau^0\big[T(G_N)\big]\Big]$, we get the convergence in distribution of traffics of G_N.

Assume now that, reciprocally, G_N has a limiting distribution of traffics τ. For any $p \geqslant 1$ and any (T,r) in $\mathcal{G}_*^p\langle \mathbf{x}\rangle$, we set

$$\begin{aligned}
\mathbb{P}_{p,N}(T,r) &= \frac{1}{\tau^0\big[(T,r)(T,r)\big]} \sum_{\substack{(H,w)\in\mathcal{G}_{*,p}\\ (H,w)\geqslant(T,r)}} \tau_N^0\big[H(G_N)\big]\mu_p\big((H,w),(T,r)\big),\\
\mathbb{P}_p(T,r) &= \frac{1}{\tau^0\big[(T,r)(T,r)\big]} \sum_{\substack{(H,w)\in\mathcal{G}_{*,p}\\ (H,w)\geqslant(T,r)}} \tau^0\big[H\big]\mu_p\big((H,w),(T,r)\big),
\end{aligned}$$

where τ^0 is the injective version of τ. By (15.2), $\mathbb{P}_p(T,r)$ is the limit of $\mathbb{P}_{p,N}(T,r)$. It belongs to $[0,1]$ and we have

$$\sum_{(T,r)\in\mathcal{G}_{*,p}} \mathbb{P}_p(T,r) = 1.$$

Hence, the collection of numbers $\mathbb{P}_p(T,r)$ well defines a random rooted graph g which is necessarily stationary. The graph g is the weak local limit of G_N.

□

16 Traffic freeness for random rooted graphs

Let $g_1 = (G_1, v_1)$ and $g_2 = (G_2, v_2)$ be two random graphs in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$. By Proposition 15.1, we can consider g_1 and g_2 as traffics in a space of traffics $(\mathcal{A}, .^*, \tau)$ on $\mathcal{G}\langle \mathbf{x}, \mathbf{x}^*\rangle$. Assume that g_1 and g_2 are free. We consider the traffic $g = g_1 + g_2$. We prove that g has the distribution of traffics of a random graph in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$, constructed inductively as below.

16.1 A free product for random rooted graphs

Let $g_1 = (G_1, v_1)$ and $g_2 = (G_2, v_2)$ be random graphs in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$. In the following, all the sampling of random graphs are independent.

1. Start by sampling to rooted graphs $g_{1,0}, g_{2,0}$ from g_1 and g_2 respectively, and identify their roots.
2. For any vertex v_1 of $g_{1,0}$ which is not its root, sample $g_{2,0v_1}$ from g_2 and identify the root of $g_{2,0v_1}$ with v_1. For any vertex v_2 of $g_{2,0}$ which is not its root, sample $g_{1,0v_2}$ from g_1 and identify the root of $g_{1,0v_2}$ with v_2.
3. For any vertex v_2 of $g_{2,0}$ which is not its root and any vertex w_1 of $g_{1,0v_2}$ which is not its root, sample $g_{2,0v_2w_1}$ from g_2 and identify the root of $g_{2,0v_2w_1}$ with w_1. And so on.

The resulting random rooted graph is called the free product of g_1 and g_2, denoted by

$$g_1 * g_2 \in \mathcal{G}_*^\infty\langle \mathbf{x}\rangle. \tag{16.1}$$

When the graphs are deterministic, this construction is also known as the free product of g_1 and g_2. Remark that $g_1 * g_2$ is uniformly bounded as soon as g_1 and g_2 are uniformly bounded.

16.2 Traffic freeness and random graphs

Proposition 16.1 (The free convolution of two random graphs)**.** *Let g_1 and g_2 in $\mathcal{G}_*^\infty\langle x\rangle$ be two integrable, traffic-stationary random graphs in one variable. Consider them in a space of traffics $(\mathcal{A}, .^*, \tau)$, being free. Then, the distribution of $g = g_1 + g_2$ is the distribution of $g_1 * g_2$.*

Proof. Let $\tilde g$ in $\mathcal{G}_*^\infty\langle x,y\rangle$ be a random graph obtained as the free product of g_1 in $\mathcal{G}_*^\infty\langle x\rangle$ and g_2 in $\mathcal{G}_*^\infty\langle y\rangle$. Denote its root by ρ. Consider a test graph $T=(V,E)$ in one variable, and chose r in V. Then, by tracking the color induced on the edges of T when injected in g_1*g_2, we obtain

$$\begin{aligned}\tau^0\big[T(g_1*g_2)\big] &= \sum_{\substack{\tilde T=(V,E,\gamma)\\ \gamma:E\to\{1,2\}}} \tau^0\big[\tilde T(\tilde g)\big]\\ &= \sum_{\substack{\tilde T=(V,E,\gamma)\\ \gamma:E\to\{1,2\}}} \mathbb{E}\Big[\mathrm{Card}\ \big\{\text{injection}(\tilde T,r)\to(\tilde g,\rho)\text{ respecting the colors.}\big\}\Big]\end{aligned}$$

Let $\tilde T$ be as in the sum. Then, by the tree like structure of $\tilde g$, two different connected components of $\tilde T$ relatively to its coloration (i.e. the partition $\pi=\big\{\{1\},\{2\}\big\}$) are always injected in independent copies of g_1 or g_2. Hence, the graph $\tilde g$ being traffic-stationary, one has

$$\tau^0\big[T(g_1*g_2)\big] \;=\; \sum_{\substack{\tilde T=(V,E,\gamma)\\ \gamma:E\to\{1,2\}}} \prod_{T_c\in\mathcal{C}_\pi(T)} \tau^0\big[T_c(g_1,g_2)\big].$$

By the definition of freeness, we get

$$\tau^0\big[T(g_1*g_2)\big] \;=\; \sum_{\substack{\tilde T=(V,E,\tilde\gamma)\\ \tilde\gamma:E\to\{1,2\}}} \tau^0\big[\tilde T(g_1,g_2)\big].$$

Hence, by formula (7.6), we get as desired $\tau^0\big[T(g_1*g_2)\big]=\tau^0\big[T(g_1+g_2)\big]$. ☐

16.3 Examples

Free product of a "Bernoulli" graph with itself: This example is interesting since it provides a rich random structure from simple objects. Let g be the random graph that is $\mathbb{Z}$ with probability p and the graph with a single vertex and no edges with probability $1-p$. Make the free product $g*g$ with itself. In Figures 4 and 18 are plotted realizations of such random graphs.

Free groups and uniform permutation matrices: A particular example of graphs are Cayley graphs of groups. Let Γ be a group. We endowed Γ with the structure of algebra of traffics $(\mathcal{A},.^*,\tau)$ in the following way. The *-algebra $(\mathcal{A},.^*)$ is the *-algebra $\mathbb{C}\Gamma$, that is the vector space of finite linear combinations

$$\sum_{\gamma\in\Gamma} a_\gamma\gamma,$$

with product

$$\sum_{\gamma\in\Gamma} a_\gamma\gamma\times\sum_{\gamma\in\Gamma} b_\gamma\gamma=\sum_{\gamma_1,\gamma_2\in\Gamma} a_{\gamma_1}b_{\gamma_2}\gamma_1\gamma_2,$$

and anti-linear involution $\gamma^*=\gamma^{-1}$. Let $\boldsymbol\gamma=(\gamma_1,\dots,\gamma_p)$ be a family of elements in Γ. We consider the Cayley graph $\Gamma_{\boldsymbol\gamma}$:

- the set of vertices of is the subgroup $\langle\boldsymbol\gamma\rangle$ of Γ spanned by $\boldsymbol\gamma$,
- for any γ in $\langle\boldsymbol\gamma\rangle$ and any $j=1,\dots,p$, vertices γ and $\gamma\gamma_j$ are linked by a directed edge with label x_j.

Rooted in the unit, $\Gamma_{\boldsymbol\gamma}$ is a graph in $\mathcal{G}_*^\infty\langle\mathbf{x}\rangle$. We set τ_γ the distribution of traffics of $\Gamma_{\boldsymbol\gamma}$. It is clear from Figure 15 that in expectation a family $(U_1^{(N)},\dots,U_p^{(N)})$ of independent uniform permutation matrices converges in distribution of traffics to $(\gamma_1,\dots,\gamma_p)$, where $\gamma_1,\dots,\gamma_p$ are the generators of the free group $\mathbb{F}_p=\langle\gamma_1,\dots,\gamma_p\rangle$.

Percolation clusters of free products of graphs: A percolation cluster of parameter $p\in[0,1]$ of a graph $g\in\mathcal{G}_*$ is a connected component of the subgraph of g obtained by removing each edge



Figure 18: A realization of the free product with itself of the random graph which is $\mathbb{Z}$ with probability 0.6, and the graph with one vertex and no edge with probability 0.4 (truncated at distance 10 of the root).

independently with probability p. We denote by $\mathfrak{C}_p(g)$ the cluster corresponding to the connected component of the root of g. This defines a random graph in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$, which is in $\mathcal{G}_*^{bnd}\langle \mathbf{x}\rangle$ as soon as g is in $\mathcal{G}_*^{bnd}\langle \mathbf{x}\rangle$.

Let g_1 and g_2 be two deterministic graphs in $\mathcal{G}_*^\infty\langle \mathbf{x}\rangle$. Then, for any $p \in [0,1]$, one has

$$\mathfrak{C}_p(g_1) * \mathfrak{C}_p(g_2) = \mathfrak{C}_p(g_1 * g_2). \tag{16.2}$$

References

[1] D. Aldous and R. Lyons. Processes on unimodular random networks. *Electron. J. Probab.*, 12:no. 54, 1454–1508, 2007.

[2] G. W. Anderson, A. Guionnet, and O. Zeitouni. *An Introduction to Random Matrices*, volume 118 of *Cambridge studies in advanced mathematics*. Cambridge University Press, 2010.

[3] I. Artemenko. Weak Convergence of Laws of Finite Graphs. *ArXiv e-prints*, March 2011.

[4] R. Basu, A. Bose, S. Ganguly, and R. Subhra Hazra. Joint convergence of several copies of different patterned random matrices. *ArXiv e-prints*, August 2011.

[5] F. Benaych-Georges and T. Lévy. A continuous semigroup of notions of independence between the classical and the free one. *Ann. Probab.*, 39(3):904–938, 2011.

[6] I. Benjamini and O. Schramm. Recurrence of distributional limits of finite planar graphs. *Electron. J. Probab.*, 6:no. 23, 13 pp. (electronic), 2001.

[7] C. Bordenave, P. Caputo, and D. Chafai. Spectrum of Markov generators on sparse random graphs. *ArXiv e-prints*, February 2012.

[8] C. Borgs, J. Chayes, L. Lovász, V. T. Sós, and K. Vesztergombi. Counting graph homomorphisms. In *Topics in discrete mathematics*, volume 26 of *Algorithms Combin.*, pages 315–371. Springer, Berlin, 2006.

[9] Włodzimierz Bryc, Amir Dembo, and Tiefeng Jiang. Spectral measure of large random Hankel, Markov and Toeplitz matrices. *Ann. Probab.*, 34(1):1–38, 2006.

[10] M. Capitaine and M. Casalis. Asymptotic freeness by generalized moments for Gaussian and Wishart matrices. Application to beta random matrices. *Indiana Univ. Math. J.*, 53(2):397–431, 2004.

[11] F. Chung. Graphlets: a Spectral Perspective of Graph Limits. *ArXiv e-prints*, March 2012.

[12] X. Ding and T. Jiang. Spectral distributions of adjacency and Laplacian matrices of random graphs. *Ann. Appl. Probab.*, 20(6):2086–2117, 2010.

[13] W. Du, X. Li, and Y. Li. The Laplacian energy of random graphs. *J. Math. Anal. Appl.*, 368(1):311–319, 2010.

[14] K. Dykema. On certain free product factors via an extended matrix model. *J. Funct. Anal.*, 112(1):31–60, 1993.

[15] Gábor Elek. On limits of finite graphs. *Combinatorica*, 27(4):503–507, 2007.

[16] A. Guionnet. *Large random matrices: lectures on macroscopic asymptotics*, volume 1957 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2009. Lectures from the 36th Probability Summer School held in Saint-Flour, 2006.

[17] L. Lovász. Very large graphs. In *Current developments in mathematics, 2008*, pages 67–128. Int. Press, Somerville, MA, 2009.

[18] L. Lovász and B. Szegedy. Limits of dense graph sequences. *J. Combin. Theory Ser. B*, 96(6):933–957, 2006.

[19] C. Male. The norm of polynomials in large random and deterministic matrices. *Probab. Theory Related Fields*, 2011.

[20] J. A. Mingo and R. Speicher. Sharp bounds for sums associated to graphs of matrices. *J.F.A.*, 262:Issue 5, p. 2272Ð2288, 2012.

[21] M. Neagu. Asymptotic freeness of random permutation matrices from gaussian matrices. *J. Ramanujan Math. Soc.*, 20(3):189–213, 2005.

[22] A. Nica. Asymptotically free families of random unitaries in symmetric groups. *Pacific J. Math.*, 157(2):295–310, 1993.

[23] A. Nica and R. Speicher. *Lectures on the combinatorics of free probability*, volume 335 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2006.

[24] D. Petz and J. Reffy. On asymptotics of large Haar distributed unitary matrices. *ArXiv Mathematics e-prints*, October 2003.

[25] D. Shlyakhtenko. Random Gaussian band matrices and freeness with amalgamation. *Internat. Math. Res. Notices*, (20):1013–1025, 1996.

[26] D. Voiculescu. Limit laws for random matrices and free products. *Invent. Math.*, 104(1):201–220, 1991.

[27] D. V. Voiculescu, K. J. Dykema, and A. Nica. *Free random variables*, volume 1 of *CRM Monograph Series*. American Mathematical Society, Providence, RI, 1992. A noncommutative probability approach to free products with applications to random matrices, operator algebras and harmonic analysis on free groups.